



**RWE Renewables UK Dogger Bank
South (East) Limited**

**RWE Renewables UK Dogger Bank
South (West) Limited**

Dogger Bank South Offshore Wind Farms


Appendix 11.1 – Marine Mammal Baseline

27/04/2023

Document Reference: 004300119

Revision: 04

Unrestricted





Company:	RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited	Asset:	Development
Project:	Dogger Bank South Offshore Wind Farms	Sub Project/Package:	Consents
Document Title or Description:	Appendix 11-1 – Marine Mammal Baseline		
Document Number:	004300119-04	Contractor Reference Number:	PC2340-RHD-OF-ZZ- RP-Z-0049

The document Originator shall complete this Cover Sheet and may give guidance below on any actions required by the recipient(s). The document Checker and Approver must not be the same person. The Document Author and Approver must not be the same person. The Approver must not be less senior than the Author.

COPYRIGHT © RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited.

All pre-existing rights reserved.

This document is supplied on and subject to the terms and conditions of the Contractual Agreement relating to this work, under which this document has been supplied, in particular:

LIABILITY

In preparation of this document RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited has made reasonable efforts to ensure that the content is accurate, up to date and complete for the purpose for which it was contracted. RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited makes no warranty as to the accuracy or completeness of material supplied by the client or their agent.

Other than any liability on RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited detailed in the contracts between the parties for this work RWE Renewables UK Dogger Bank South (West) Limited and RWE Renewables UK Dogger Bank South (East) Limited shall have no liability for any loss, damage, injury, claim, expense, cost or other consequence arising as a result of use or reliance upon any information contained in or omitted from this document.

Any persons intending to use this document should satisfy themselves as to its applicability for their intended purpose.

The user of this document has the obligation to employ safe working practices for any activities referred to and to adopt specific practices appropriate to local conditions.

Rev No.	Date	Status/Reason for Issue	Author	Checked by	Approved by
01	15/03/2023	First Issue	AS/SB	HC	AP
02	04/04/2023	Second Issue	SB	AS	HC
03	24/04/2023	Third Issue	LL	AS	HC
04	27/04/2023	Final Issue	LL	CC	AC

Unrestricted

Contents

11	Marine Mammal Information.....	6
11.1	Introduction.....	6
11.2	Impact Scoping	6
11.3	Marine Mammal Species	7
11.4	Study Area.....	9
11.4.1	Policy, Legislation and Guidance	10
11.4.1.1	National Policy Statements	10
11.4.1.3	The Marine Strategy Framework Directive	11
11.4.1.4	The Marine Policy Statement.....	11
11.4.1.5	The East Inshore and East Offshore Marine Plans	12
11.4.2	Other National and International Legislation for Marine Mammals	12
11.4.3	European Protected Species guidance	17
11.4.3.2	Marine Wildlife Licence Requirements	19
11.4.3.3	Conservation Status of Marine Mammals.....	19
11.5	Site Specific Surveys	21
11.5.1	DBS Site Specific Surveys.....	21
11.5.2	Surveys Within the Dogger Bank Area.....	27
11.5.4	Other Available Sources.....	31
11.6	Existing Environment	33
11.6.1	Harbour Porpoise.....	33
11.6.1.1	Desk-Based Review of Harbour Porpoise Presence.....	33
11.6.1.2	Results from the Site-Specific Surveys for Harbour Porpoise.....	35
11.6.1.3	Results from Surveys in the Dogger Bank Area.....	35
11.6.1.4	Site-Specific Density Estimates for Harbour Porpoise	35
11.6.1.5	Review of Abundance and Density Estimates for Harbour Porpoise	42
11.6.1.6	Diet of Harbour Porpoise	45
11.6.2	Bottlenose Dolphin	46
11.6.2.1	Desk-Based Review of Bottlenose Dolphin Presence	46
11.6.2.2	Results from the Site-Specific Surveys for Bottlenose Dolphin	50
11.6.2.3	Results from Surveys in the Dogger Bank Area.....	50
11.6.2.4	Site-Specific Density Estimates for Bottlenose dolphin	50

11.6.2.5	Review of Abundance and Density Estimates for Bottlenose Dolphins.....	50
11.6.2.6	Diet of Bottlenose Dolphin.....	52
11.6.3	Common Dolphin	53
11.6.3.1	Desk-Based Review of Common Dolphin Presence	53
11.6.3.2	Results from the Site-Specific Surveys for Common Dolphin	55
11.6.3.3	Results from Surveys in the Dogger Bank Area.....	56
11.6.3.4	Site-Specific Density Estimates for Common Dolphin.....	56
11.6.3.5	Review of Abundance and Density Estimates for Common Dolphin	57
11.6.3.6	Diet of Common Dolphin.....	58
11.6.4	White-Beaked Dolphin	59
11.6.4.1	Desk-Based Review of White-Beaked Dolphin Presence	59
11.6.4.2	Results from the Site-Specific Surveys for White-Beaked Dolphin.....	60
11.6.4.3	Results from Surveys in the Dogger Bank Area.....	60
11.6.4.4	Site-Specific Density Estimates for White-Beaked Dolphin	61
11.6.4.5	Review of Abundance and Density Estimates for White-Beaked Dolphin.....	61
11.6.4.6	Diet of White-Beaked Dolphin	62
11.6.5	Minke Whale.....	63
11.6.5.1	Desk-Based Review of Minke Whale Presence.....	63
11.6.5.2	Results from the Site-Specific Surveys for Minke Whale	65
11.6.5.3	Results from Surveys in the Dogger Bank Area.....	65
11.6.5.4	Site-Specific Density Estimates for Minke Whale.....	66
11.6.5.5	Review of Abundance and Density Estimates for Minke Whale	66
11.6.5.6	Diet of Minke whale.....	68
11.6.6	Grey Seal	68
11.6.6.1	Desk-Based Review of Grey Seal Presence	68
11.6.6.2	Grey Seal Haul-Out Sites.....	71
11.6.6.3	Results from the Site-Specific Surveys for Grey Seal	75
11.6.6.4	Site-Specific Density Estimates for Grey Seal.....	76
11.6.6.5	Review of Abundance and Density Estimates for Grey Seal.....	81
11.6.6.6	Diet and Foraging of Grey Seal	84
11.6.7	Harbour seal	85
11.6.7.1	Desk-Based Review of Harbour Seal Presence.....	85
11.6.7.2	Harbour Seal Haul-Out Sites.....	89

11.6.7.3	Results from the Site-Specific Surveys for Harbour Seals	91
11.6.7.4	Results from Surveys in the Dogger Bank Area.....	91
11.6.7.5	Site-Specific Density Estimates for Harbour Seal.....	91
11.6.7.6	Review of Abundance and Density Estimates for Harbour Seal.....	92
11.6.7.7	Diet and Foraging of Harbour Seal	94

Tables

Table 11-1	Summary Table for National and International Legislations Relevant for Marine Mammals	13
Table 11-2	Conservation Status of Marine Mammal Species Occurring in UK and Adjacent Waters (JNCC 2019), Relevant to the Project	20
Table 11-3	Global IUCN Red List of Threatened Species Assessments for Marine Mammal Species Relevant to the Project.....	20
Table 11-4	APEM Surveys Species Counts for DBS East Array Area and 4km Buffer (March 2021 to February 2022).....	25
Table 11-5	APEM Surveys Species Counts for DBS West Array Area and 4km Buffer (March 2021 to February 2022).....	26
Table 11-6	Other Available Data and Information Sources.....	31
Table 11-7	Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km ²) of Harbour Porpoises in: a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer	36
Table 11-8	Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km ²) of Harbour Porpoises in: a) DBS West Array Area plus 4km buffer; b) DBS West Array Area c) 4km buffer	37
Table 11-9	Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km ²) of Unidentified Dolphin/Porpoise in a) DBS East Array Area plus 4km Buffer; b) DBS East Array Area c) 4km Buffer.....	39
Table 11-10	Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km ²) of Unidentified Dolphin/Porpoise in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer	40
Table 11-11	Seasonal Density Estimates for Harbour Porpoise from APEM Ltd Survey	42
Table 11-12	Seasonal Density Estimates for Bottlenose Dolphin from Waggitt et al. (2019) Densities	51
Table 11-13	Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km ²) of Common Dolphins: a) DBS West Array Area Plus 4km; b) DBS West Array Area c) 4km Buffer	56
Table 11-14	Seasonal Density Estimates for Common Dolphin from Waggitt et al. 2019 Densities	57

Table 11-15 Seasonal Density Estimates for White-beaked Dolphin from Waggitt et al. 2019 Densities.....	62
Table 11-16 The Most Recent Grey Seal Count at Each of the Nearby Haul-out Sites, and the Distance to DBS	72
Table 11-17 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km ²) of Grey Seal in: a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer.....	76
Table 11-18 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km ²) of Grey Seal in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer.....	77
Table 11-19 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km ²) of Unidentified Seal Species in a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer	78
Table 11-20 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km ²) of Unidentified Seal Species in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer	79
Table 11-21 Grey Seal Count Population Estimates	84
Table 11-22 The Most Recent Harbour Seal Count at Each of the Nearby Haul-out Sites, and the Distance to DBS.....	89
Table 11-23 Harbour Seal Counts and Population Estimates in the SE England MU.....	94

Plates

Plate 11-1 Tagged Grey Seal Movements Along the East Coast of England (Carter et al. 2020).	7
Plate 11-2 Survey Area for DBS East Array Area with 4km Buffer (Including Extension) and 2km Transects.....	22
Plate 11-3 Survey area for DBS West array area with 4km buffer (including extension) and 2km transects.....	23
Plate 11-4 Location of Aerial Survey Transects Within Dogger Bank and Tranche C.	28
Plate 11-5 Boat-based Survey Transect Lines in Dogger Bank (Gardline).....	29
Plate 11-6 Incidental Odontocete Sightings During Ornithology Surveys for the Forewind Dogger Bank Offshore Wind Farm. January 2010 to June 2012	30
Plate 11-7 Spatial Variation in Predicted Densities (Individuals per km of Harbour Porpoise in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.....	34
Plate 11-8 Estimated Abundance of Harbour Porpoise Within DBS East Array Area.....	41
Plate 11-9 Estimated Abundance of Harbour Porpoise Within DBS West Array Area	41

Plate 11-10 Area Covered by the SCANS-III Survey, and the Locations of the SCANS-III Survey Blocks (The Projects Are Within Survey Block O) (Hammond et al. 2021).....	43
Plate 11-11 Estimated Harbour Porpoise Density in Each SCANS-III Survey Block (Hammond et al. 2021).....	44
Plate 11-12 Harbour porpoise Management Units (MUs), noting that this species is largely confined to the continental shelf (i.e. waters <200m depth) (IAMMWG, 2022).....	45
Plate 11-13 Estimated Bottlenose Dolphin Density in Each SCANS-III Survey Block (Hammond et al. 2021).....	48
Plate 11-14 Spatial Variation in Predicted Densities (Individuals per km of Bottlenose Dolphin in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.....	49
Plate 11-15 Bottlenose Dolphin MU (IAMMWG, 2022).....	52
Plate 11-16 Estimated Common Dolphin Density in Each SCANS-III Survey Block (Hammond et al. 2021).....	54
Plate 11-17 Spatial Variation in Predicted Densities Individuals per km of Common Dolphin in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.....	55
Plate 11-18 Common Dolphin, White-beaked Dolphin and Minke Whale MU (IAMMWG, 2022).....	58
Plate 11-19 Spatial Variation in Predicted Densities Individuals per km of White-beaked Dolphin in January and July in the NE Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.....	60
Plate 11-20 Spatial Variation in Predicted Densities (Individuals per km of Minke Whale in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.....	65
Plate 11-21 Estimated Minke Whale Density in Each SCANS-III Survey Block (Hammond et al. 2021)	67
Plate 11-22 Tagged Grey Seal Movements Along the East Coast of England (Russell 2016).	70
Plate 11-23 Grey Seal Telemetry Tags (Shown in Green are the Results from the Tagging of 11 Individuals in 2012, from the Baie de Somme (BDS) Haul-out Site on the North Coast of France. Red Dots Indicate Haul-out Sites, and Red Circles Indicate Breeding Locations.....	71
Plate 11-24 Estimated Abundance of Grey Seal at DBS East Array Area.....	80
Plate 11-25 Estimated Abundance of Grey Seal at DBS West Array Area.....	80
Plate 11-26 Relative Density of Grey Seals a) Mean With Associated Cell-wise b) Lower and c) Upper 95% Confidence Intervals of the Percentage of the at Sea Population (Excluding Hauled-out Seals) Estimated to be Present in Each 5km x 5km Grid Cell at Any One Time (Carter et al. 2022).....	82

Plate 11-27 Telemetry Tracks by Deployment Region for Harbour Seals Aged One Year or Over (Russell and McConnell 2014).....86

Plate 11-28 Selected Examples of Foraging Trips. Purple: Elsam02 On an 18-day Trip to the Deeper Parts of the North Sea.. Red: Elsam05 on a 15-day Trip to the German Bight. Blue: Elsam08 on a 10-day Trip to Horns Reef (Tougaard et al. 2003).....88

Plate 11-29 Relative Density of Harbour Seals a) Mean With Associated Cell-wise b) Lower and c) Upper 95% Confidence Intervals of the Percentage of the at Sea Population (Excluding Hauled-out Seals) Estimated to be Present in Each 5km x 5km Grid Cell at Any One Time.92

Figures

Figure 11-1-1 Grey Seal At-sea Distribution. Maps Show Mean Percentage of At-sea Population Estimated to be Present in Each 5km x 5 km Grid Square at Any One Time, and the Square-wise (Carter et al. 2022)

Figure 11-1-2 Harbour Seal at Sea Distribution. Maps Show Mean Percentage of At-sea Population Estimated to be Present in Each 5km x 5km Grid Square at Any One Time, and the Square-wise (Carter et al. 2022)

Glossary

Term	Definition
Array Areas	The DBS East and DBS West offshore array areas, where the wind turbines and array cables will be located. The array areas do not include the offshore export cable corridor.
Cetaceans	Commonly known as whales, dolphins, or porpoises.
Dogger Bank South (DBS) Offshore Wind Farms	The collective name for the two Projects, DBS East and DBS West.
European Sites	Sites designated for nature conservation under the Habitats Directive and Birds Directive. This includes candidate Special Areas of Conservation, Sites of Community Importance, Special Areas of Conservation and Special Protection Areas, and is defined in regulation 8 of the Conservation of Habitats and Species Regulations 2017.
Landfall	The point on the coastline at which the offshore export cables are brought onshore, connecting to the onshore export cables at the Transition Joint Bay.
Management Unit	Management units provide an indication of the spatial scales at which impacts of plans and projects alone, cumulatively and in-combination, need to be assessed for the key cetacean species in UK waters, with consistency across the UK.
Offshore Development Area	The Offshore Development Area encompasses the wind farm site and the options for the offshore transmission works.
Offshore export Cable Corridor	This is the area which includes cable trenches, haul roads, spoil storage areas, and limits of deviation for micro-siting. For assessment purposes, the cable corridor does not include the onshore substations, Transition Joint Bays or temporary access routes; but includes construction compounds (purely for the cable route).

Term	Definition
Pinnipeds	Commonly known as seals.
Projects	DBS East and DBS West (collectively referred to as the Dogger Bank South offshore wind farms).

Acronyms

Term	Definition
ASCOBANS	Agreement on the Conservation of Small Cetaceans of the Baltic, North East Atlantic, Irish and North Seas
BDS	Baie de Somme
BEIS	Department for Business, Energy & Industrial Strategy
CCW	Countryside Council for Wales
CES	Coastal East Scotland
CI	Confidence Intervals
CIS	Celtic and Irish Sea
CITES	Convention on International Trade in Endangered Species of Wild Fauna and Flora
CGNS	Celtic and Greater North Sea
COWRIE	Collaborative Offshore Wind Research into the Environment
CroW	The Countryside and Rights of Way Act
CV	Coefficient of Variation
DBS	Dogger Bank South
DECC	Department of Energy and Climate Change
EC	European Communities
EPS	European Protected Species
ESAS	European Seabirds At Sea
FCS	Favourable Conservation Status
GNS	Greater North sea

Term	Definition
HiDef	HiDef Aerial Surveying Limited
HM	His Majesty's
IAMMWG	Inter-Agency Marine Mammal Working Group
IUCN	International Union for Conservation of Nature
JCP	Joint Cetacean Protocol
JNCC	Joint Nature Conservation Committee
km	Kilometre
MMO	Marine Management Organisation
MMMP	Marine Mammal Mitigation Protocol
MPS	Marine Policy Statement
MSFD	Marine Strategy Framework Directive
MUs	Management Units
NE	North East
NPS	National Policy Statements
NS	North Sea
NSIP	Nationally Significant Infrastructure Project
OESEA	Offshore Energy Strategic Environmental Assessment
ORCA	Organisation Cetacea
OSPAR	Oslo and Paris Convention for the Protection of the Marine Environment 1992
PAMS	Passive Acoustic Monitoring System

Term	Definition
QC	Quality Control
SAC	Special Area of Conservation
SCANS	Small Cetaceans in European Atlantic waters and the North Sea
SCOS	Special Committee on Seals
SE	South East
TSEG	Trilateral Seal Expert Group
UK	United Kingdom
WS	West Scotland
WWT	The Wildfowl and Wetlands Trust

11 Marine Mammal Information

11.1 Introduction

1. This appendix provides further supporting marine mammal information and survey data for **Chapter 11 Marine Mammal Ecology**.

11.2 Impact Scoping

2. The potential for impact from physical barrier effects during operation has been scoped out.
3. Monitoring studies at Nysted and Rødsand have indicated that operational activities have had no impact on regional seal populations (Teilmann *et al.* 2006; McConnell *et al.* 2012). Tagged harbour seals have been recorded within two operational offshore wind farm sites (Alpha Ventus in Germany and Sheringham Shoal in UK) with the movement of several of the seals suggesting foraging behaviour around wind turbines (Russell *et al.* 2014). Both harbour porpoise and seals have been shown to forage within operational offshore wind farms (e.g. Lindeboom *et al.* 2011; Russell *et al.* 2014), indicating no restriction to movements in operational offshore wind farm sites.
4. **Plate 11-1** shows tagged grey seal movements around the UK coastlines from 114 grey seal (left) and 239 harbour seals (right). These tagging studies indicate that grey seal associated with haul-out sites on the east coast of England forage at significant distances offshore, with grey seals travelling through the Offshore Development Area (Carter *et al.* 2020). For harbour seal, the tagging studies show a smaller foraging range than for grey seal, with limited potential for connectivity with the Offshore Development Area. However, as noted above, seals are known to still utilise operational wind farm areas, and there is no indication that the physical structures would cause a barrier to their movement, or a reduction in their foraging.
5. Effects on harbour porpoise are more difficult to assess as various operational activities may influence the species differently. Teilman and Carstensen (2012) have found that harbour porpoise may habituate itself to the wind farm post-construction (possibly due to habitat enrichment and reduced fishing) but the physical presence of the turbines is unlikely to create a barrier to the species (Tougaard *et al.* 2005).
6. The spacing of 830m between wind turbines would allow animals to move between infrastructure and through the operational wind farm site. This means that animals can be expected to move within and through the operational wind farm, irrespective of layout.




Plate 11-1 Tagged Grey Seal Movements Along the East Coast of England (Carter et al. 2020).

7. Based on the limited potential for any disturbance (or barrier to movement) due to the presence of the wind farm infrastructure, and that the spacing would allow for marine mammals to transit through the wind farm site while maintaining distance between themselves and the infrastructure, it is not anticipated that there would be any potential for a barrier to marine mammal movement, and this potential effect has been scoped out of further assessment.

11.3 Marine Mammal Species

8. In United Kingdom (UK) waters, two groups of marine mammals occur: cetaceans (whales, dolphins, and porpoises) and pinnipeds (seals). During the site specific surveys for both the Dogger Bank South (DBS) offshore wind farms, harbour porpoise *Phocoena phocoena* were the most commonly sighted marine mammal species for both Projects, with the highest numbers being recorded in the summer months) (APEM Ltd, 2022).
9. This is supported by other wider scale surveys and reporting for marine mammals in the area, including by Department for Energy and Climate Change (DECC) (2016), Small Cetaceans in European Atlantic waters and the North Sea (SCANS) surveys (Hammond et al. 2017) and Joint Cetacean Protocol (JCP) data resources (Paxton et al. 2016).

10. While a number of cetacean species have been recorded within the southern areas of the North Sea, only harbour porpoise occur regularly throughout the year, while minke whale *Balaenoptera acutorostrata* could occur in the area, particularly during in the summer periods and white-beaked dolphin *Lagenorhynchus albirostris* are less frequent (DECC 2016; Hammond *et al.* 2017; Paxton *et al.* 2016). Other cetacean species, including common dolphin *Delphinus delphis*, bottlenose dolphin *Tursiops truncatus* and white-sided dolphin *Lagenorhynchus acutus* are relatively uncommon in the area (DECC 2016). Although it should be noted that the number of bottlenose dolphin sightings has recently been increasing along the east coast of England, where a pod has been announced to be resident off the coast of Yorkshire (The Telegraph 2022).
11. Both UK seal species, grey seal, *Halichoerus grypus*, and harbour seal, *Phoca vitulina*, are present in the area in relatively high number, due to nearby key breeding areas for both species (DECC 2016).
12. The most recent public sightings reported to the SeaWatch Foundation in the east of England (at the time of writing; February 2022 to January 2023) were predominantly harbour porpoise 59 sightings (n=133), bottlenose dolphin; 71 sightings (n=964), white-beaked dolphin; two sightings (n=5), common dolphin five sightings (n=61), minke whale; four sightings (n=4), humpback whale; one sightings (n=2), unknown cetacean species (n=1), grey seal; 38 sightings (n=94), and harbour seal (n=1), (SeaWatch Foundation 2023).
13. Other marine mammal species, including Atlantic white-sided dolphin, bottlenose dolphin, killer whale, *Orcinus orca*, sperm whale, *Physeter macrocephalus*, long-finned pilot whale, *Globicephala melas*, Risso's dolphin, *Grampus griseus*, striped dolphin, *Stenella coeruleoalba*, and other seal species are occasional or rare visitors to the southern North Sea (e.g. Reid *et al.* 2003; Hammond *et al.* 2013, 2017; DECC 2016; SCOS 2020).
14. Sperm whale are a regular visitor to the Norwegian Sea and have been recorded to strand in the southern North Sea (Seawatch Foundation, 2016). In recent years there have been sightings of walrus (*Odobenus rosmarus*) in the North Sea, with the most recent being recorded in December 2022 in Scarborough (The Guardian, 2023).
15. Site characterisation has been undertaken using site specific data for DBS, as well as existing data from other Dogger Bank offshore wind farms and other available information for the region.

16. The study area for marine mammals has been defined on the basis of marine mammals being highly mobile and transitory in nature; therefore, it is necessary to examine species occurrence not only within the DBS Project areas, but also over the wider area. Based on the site-specific surveys and other data sources, the key species of interest and therefore the focus of the assessments will be on the following species:
- Harbour porpoise – present throughout the year, although may be variations in seasonal occurrence;
 - Bottlenose dolphin – historically not common in the area, with limited data, however, recent reporting has indicated that the number are increasing in the area, and so have been included on a precautionary basis;
 - Common dolphin – seasonal occurrence in low numbers but historically not common in the area;
 - White-beaked dolphin – seasonal occurrence in low numbers;
 - Minke whale – seasonal occurrence in low numbers;
 - Grey seal – present throughout the year; and
 - Harbour seal – present throughout the year.

11.4 Study Area

17. Management Units (MUs) provide an indication of the spatial scales at which effects of plans and projects alone, and in-combination, need to be assessed for the key cetacean species in UK waters with consistency across the UK (IAMMWG 2022). The study area, MUs and reference populations have been determined based on the most relevant information and scale at which potential impacts from DBS wind farms alone.
18. For each species of marine mammal, the following study areas have been defined based on the relevant MUs, current knowledge and understanding of the biology of each species:
- Harbour porpoise: North Sea (NS) MU;
 - Bottlenose dolphin: Greater North Seas (GNS) MU;
 - Common Dolphin: Celtic and Greater North Seas (CGNS) MU;
 - White beaked dolphin: CGNS MU;
 - Minke whale: CGNS MU;
 - Grey seal: South-east (SE) England, and the North-east England (NE) MUs; and
 - Harbour seal: SE England MU.

19. There is the potential for seals from haul-out sites to move along the coast and offshore to forage in and around the proposed Project areas. Key haul-out sites for both seal species within the vicinity of the DBS Projects are:
- Filey Brigg;
 - Ravenscar;
 - Tees;
 - The Wash;
 - Donna Nook; and
 - Blakeney Point.

11.4.1 Policy, Legislation and Guidance

11.4.1.1 National Policy Statements

20. As outlined in the **Chapter 11 Marine Mammal Ecology** there are a number of pieces of legislation, policy and guidance applicable to the assessment of marine mammals. These include:
- The Marine Strategy Framework Directive (MSFD) 2008/56/EC (European Communities 2008);
 - The Marine Policy Statement (MPS) (His Majesty's (HM) Government 2011); and
 - The East Inshore and East Offshore Marine Plans (HM Government, 2014).

11.4.1.3 The Marine Strategy Framework Directive

21. Annex I of the MSFD states that to ensure that good environmental status is met the following must be considered:
- Biological diversity should be maintained;
 - The quality and occurrence of habitats, as well as the distribution and abundance of species are in line with prevailing physiographic, geographic, and climatic conditions;
 - All elements of the marine food web, to the extent that they are known, occur at normal abundance and diversity levels capable of ensuring the long-term abundance of the species and the retention of their full reproductive capacity;
 - Concentrations of contaminants are at levels not giving rise to pollution effects;
 - Properties and quantities of marine litter do not cause harm to the coastal and marine environment; and
 - Introduction of energy, including underwater noise, is at levels that do not adversely affect the marine environment.

11.4.1.4 The Marine Policy Statement

22. The MPS (HM Government 2011) provides a high-level approach to marine planning and the general principles for decision making. It sets out the framework for environmental, social and economic considerations that need to be taken into account in marine planning. The high-level objective of 'Living within environmental limits' covers the points relevant to marine mammals, this requires that:
- Biodiversity is protected, conserved and where appropriate recovered and loss has been halted;
 - Healthy marine and coastal habitats occur across their natural range and are able to support strong, biodiverse biological communities and the functioning of healthy, resilient and adaptable marine ecosystems; and
 - Our oceans support viable populations of representative, rare, vulnerable, and valued species.

11.4.1.5 The East Inshore and East Offshore Marine Plans

23. Within both the East Inshore and East Offshore Marine Plans (HM Government 2014), a set of objectives have been set out to ensure biodiversity protections and are of relevance to marine mammals as they cover policies and commitments on the wider ecosystem, as set out within the MPS and the MSFD.

- Objective 6: “To have a healthy, resilient and adaptable marine ecosystem in the East Marine Plan areas”; and
- Objective 7: “To protect, conserve and, where appropriate, recover biodiversity that is in or dependent upon the East marine plan areas”.

11.4.2 Other National and International Legislation for Marine Mammals

24. **Table 11-1** provides an overview of national and international legislation in relation to marine mammals.

Table 11-1 Summary Table for National and International Legislations Relevant for Marine Mammals

Legislation	Level of Protection	Species Included	Details
Agreement on the Conservation of Small Cetaceans of the Baltic, North East Atlantic, Irish and North Seas (ASCOBANS)	International	Odontocetes	Formulated in 1992, this agreement has been signed by eight European countries bordering the Baltic and North Seas (including the English Channel) and includes the UK. Under the Agreement, provision is made for the protection of specific areas, monitoring, research, information exchange, pollution control and increasing public awareness of small cetaceans.
The Berne Convention 1979	International	All cetaceans, grey seal and harbour seal	The Convention conveys special protection to those species that are vulnerable or endangered. Appendix II (strictly protected fauna): 19 species of cetacean. Appendix III (protected fauna): all remaining cetaceans, grey and harbour seal. Although an international convention, it is implemented within the UK through the Wildlife and Countryside Act 1981 (with any aspects not implemented via that route brought in by the Habitats Directive).
The Bonn Convention 1979	International	All cetaceans	Protects migratory wild animals across all, or part of their natural range, through international co-operation, and relates particularly to those species in danger of extinction. One of the measures identified is the adoption of legally binding agreements, including ASCOBANS.

Legislation	Level of Protection	Species Included	Details
Oslo and Paris Convention for the Protection of the Marine Environment 1992 (OSPAR)	International	Bowhead whale <i>Balaena mysticetus</i> , northern right whale <i>Eubalaena glacialis</i> , blue whale <i>Balaenoptera musculus</i> , and harbour porpoise	OSPAR has established a list of threatened and/or declining species in the North East Atlantic. These species have been targeted as part of further work on the conservation and protection of marine biodiversity under Annex V of the OSPAR Convention. The list seeks to complement, but not duplicate, the work under the EC Habitats and Birds directives and measures under the Berne Convention and the Bonn Convention.
International Convention for the Regulation of Whaling 1956	International	All cetacean species	This Convention established the International Whaling Commission (IWC) who regulates the direct exploitation and conservation of large whales (in particular sperm and large baleen whales) as a resource and the impact of human activities on cetaceans. The regulation considered scientific matters related to small cetaceans, in particular the enforcing of a moratorium on commercial whaling which came into force in 1986.

Legislation	Level of Protection	Species Included	Details
Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) 1975	International	All cetacean species	Prohibits the international trade in species listed in Annex 1 (including sperm whales, northern right whales, and baleen whales) and allows for the controlled trade of all other cetacean species.
The Conservation of Habitats and Species Regulations 2017 and The Conservation of Offshore Marine Habitats and Species Regulations 2017	National	All cetaceans, grey and harbour seal	The Habitats Regulations 2017. Provisions of The Habitats Regulations are described further in Chapter 11 Marine Mammals. It should be noted that the Habitats Regulations apply within the territorial seas and to marine areas within UK jurisdiction, beyond 12 nautical miles.

Legislation	Level of Protection	Species Included	Details
The Wildlife and Countryside Act 1981 (as amended)	National	All cetaceans	Schedule five: all cetaceans are fully protected within UK territorial waters. This protects them from killing or injury, sale, destruction of a particular habitat (which they use for protection or shelter) and disturbance. Schedule six: Short-beaked common dolphin, bottlenose dolphin and harbour porpoise; prevents these species being used as a decoy to attract other animals. This schedule also prohibits the use of vehicles to take or drive them, prevents nets, traps or electrical devices from being set in such a way that would injure them and prevents the use of nets or sounds to trap or snare them.
The Countryside and Rights of Way Act (CroW) 2000	National	All cetaceans	Under the CroW Act 2000, it is an offence to intentionally or recklessly disturb any wild animal included under Schedule 5 of the Wildlife and Countryside Act.
Conservation of Seals Act 1970	International	Grey and harbour seal	As of 1st March 2021, a person commits an offence if they intentionally or recklessly kill, injure or take a seal. The legislative changes in England and Wales, amends the Conservation of Seals Act 1970, prohibiting the intentional or reckless killing, injuring or taking of seals and removing the provision to grant licences for the purposes of protection, promotion or development of commercial fisheries or aquaculture activities. These changes were enacted to ensure compliance with the US Marine Mammal Protection Act Import Provision Rule.

11.4.3 European Protected Species guidance

25. All cetacean species are listed as European Protected Species (EPS) under Annex IV of The Habitats Directive and are therefore protected from the deliberate killing (or injury), capture and disturbance throughout their range. Within the UK, The Habitats Directive is enacted through The Conservation of Habitats and Species Regulations 2017 and the Conservation of Offshore Marine Habitats and Species Regulations 2017. Under these Regulations, it is an offence if cetaceans are deliberately disturbed in such a way as to:
 - Deliberately capture, injure or kill any EPS;
 - Deliberately disturb them; or
 - Damage or destroy a breeding site or resting place.
26. The Joint Nature Conservation Committee (JNCC), Natural England and the Countryside Council for Wales (CCW)¹ (JNCC *et al.* 2010) produced draft guidance concerning the Regulations on the deliberate disturbance of marine EPS, which provides an interpretation of the regulations in greater detail, including for pile driving operations (JNCC 2010a), seismic surveys (JNCC 2017) and the use of explosives (JNCC 2010b).
27. The draft guidance provides the following interpretations of deliberate injury and disturbance offences under both the Habitats Regulations and Offshore Regulations (now the Habitats Regulations 2017), as detailed in the paragraphs below:

“Deliberate actions are to be understood as actions by a person who knows, in light of the relevant legislation that applies to the species involved, and the general information delivered to the public, that his action will most likely lead to an offence against a species, but intends this offence or, if not, consciously accepts the foreseeable results of his action;

Certain activities that produce loud sounds in areas where EPS could be present have the potential to result in an injury offence, unless appropriate mitigation measures are implemented to prevent the exposure of animals to sound levels capable of causing injury”.

¹ Now part of Natural Resources Wales

29. For the purposes of marine users, the draft guidance states that a disturbance which can cause offence should be interpreted as:
“Disturbance which is significant in that it is likely to be detrimental to the animals of an EPS or significantly affect their local abundance or distribution”.
30. The draft guidelines further states that a disturbance offence is more likely where an activity causes persistent noise in an area for long periods of time, and highlights that sporadic “trivial disturbance” should not be considered as a disturbance offence under Article 12.
31. Any action that could increase the risk of a long-term decline of the population, increase the risk of a reduction of the range of the species, and/or increase the risk of a reduction of the size of the habitat of the species can be regarded as a disturbance under the Regulations. For a disturbance to be considered non-trivial, the disturbance to marine EPS would need to be likely to at least increase the risk of a certain negative impact on the species at Favourable Conservation Status (FCS).
32. JNCC *et al.* (2010) state that:
“In any population with a positive rate of growth, or a population remaining stable at what is assumed to be the environmental carrying capacity, a certain number of animals can potentially be removed as a consequence of anthropogenic activities (e.g., through killing, injury or permanent loss of reproductive ability), in addition to natural mortality, without causing the population to decrease in numbers, or preventing recovery, if the population is depleted. Beyond a certain threshold however, there could be a detrimental effect on the population”.
33. Further discussion on the use of thresholds for significance and the permanent or temporary nature of any disturbance is considered by defining the magnitude of potential effect in the assessment. Consideration of any potential essential habitat or geographical structuring of EPS is provided in the existing environment section.

11.4.3.2 Marine Wildlife Licence Requirements

34. A marine wildlife licence is required if the risk of injury or disturbance to cetacean species is assessed as likely under the Habitats Regulations 2017. If a licence is required, an application must be submitted, the assessment of which comprises three tests, namely:
 - Whether the activity falls within one of the purposes specified in Regulation 55 of the Habitats Regulations. Only the purpose of “*preserving public health or public safety or other imperative reasons of overriding public interest, including those of a social or economic nature and beneficial consequences of primary importance for the environment*” is of relevance to marine mammals in this context;
 - That there are no satisfactory alternatives to the activity proposed (that would not incur the risk of offence); and
 - That the licensing of the activity will not result in a negative impact on the species’/ population’s FCS.
35. A marine wildlife licence would consider all cetacean species at potential risk of injury or disturbance. There is no legislation that requires seals to be included under a marine wildlife licence; disturbance is not an offence under the Conservation of Seals Act 1970 and, in the case of injury to seals, the Marine Management Organisation (MMO) is only able to grant licences under very specific circumstances as listed under Section 10(1) of the Conservation of Seals Act 1970 which would not apply in the case that a marine wildlife licence was required for the construction of DBS arrays.
36. Under the definitions of ‘deliberate disturbance’ in the Habitats Regulations, chronic exposure and / or displacement of animals could be regarded as a disturbance offence. Therefore, if these risks cannot be avoided, then the Applicant is likely to be required to apply for a marine wildlife licence from the MMO in order to be exempt from the offence.
37. If required, the marine wildlife licence application will be submitted post-consent. At that point in time, the project design envelope will have been further refined through detailed design and procurement activities and further detail will be available on the techniques selected for the construction of the wind farm, as well as the mitigation measures that will be in place following the development of the Marine Mammal Mitigation Protocol (MMMP) for piling.

11.4.3.3 Conservation Status of Marine Mammals

38. **Table 11-2** provides the current conservation status of marine mammal species occurring in UK and adjacent waters, based on the most recent 2013-2018 reporting by JNCC in 2019.

Table 11-2 Conservation Status of Marine Mammal Species Occurring in UK and Adjacent Waters (JNCC 2019), Relevant to the Project

Species	Conservation Status Assessment
Harbour porpoise	Unknown
Bottlenose dolphin	Unknown
White-beaked dolphin	Unknown
Minke whale	Unknown
Grey seal	Favourable
Harbour seal	Unfavourable-inadequate

39. The International Union for Conservation of Nature (IUCN)'s Red List of Threatened Species provides assessments of the conservation status of animals evaluated at a global scale using the IUCN Red List Categories and Criteria, with the aim of determining their relative risk of extinction. Assessments are updated periodically to reflect new information. Where sufficient information exists, the majority of marine mammal species occurring in UK waters fall into the lowest category of 'least concern' (**Table 11-3**).

Table 11-3 Global IUCN Red List of Threatened Species Assessments for Marine Mammal Species Relevant to the Project

Species	IUCN Red List Status	Year Assessed
Harbour porpoise	Least Concern	2020
Bottlenose dolphin	Least Concern	2018
White-beaked dolphin	Least Concern	2018
Minke whale	Least Concern	2018
Grey seal	Least Concern	2016
Harbour seal	Least Concern	2016

11.5 Site Specific Surveys

11.5.1 DBS Site Specific Surveys

40. In order to provide site specific and up to date information on which to base the impact assessment, a site-specific digital aerial survey campaign was conducted for both marine mammals and seabirds. APEM Limited ('APEM') collected high resolution digital aerial still imagery for marine megafauna (combined with ornithology surveys) over both DBS East and DBS West, including a 4km buffer (see **Plate 11-2** and **Plate 11-3**).
41. Three standard APEM cameras with sensors set to a resolution of 1.5cm Ground Sample Distance were used.
42. The digital aerial survey was a grid-based design conducted along a series of strip transects (at 2km spacing), flown on a monthly basis from March 2021 to February 2022. The camera system captured abutting still imagery along survey transit lines spaced approximately 2km between-track (see **Plate 11-2** and **Plate 11-3**)
43. The aircraft collected the data at an altitude of approximately 396m and a speed of approximately 120 knots. Images were collected continuously along the survey lines and a minimum of 30% coverage was captured, with 10% of the data subject to further analysis.



Plate 11-2 Survey Area for DBS East Array Area with 4km Buffer (Including Extension) and 2km Transects



Plate 11-3 Survey area for DBS West array area with 4km buffer (including extension) and 2km transects

44. Imagery was captured in raw format and post-processed to ensure optimal quality for the subsequent stage of image analysis, to extract information on marine fauna or other notable occurrences.
45. Data analysis follows a two-stage process in which images are reviewed (10%) then the detected objects are identified to species or species group level. The images undergo quality control inhouse.
46. Density and abundance estimates are calculated using the raw counts divided by the number of images collected to give the mean number of animals per image. Population estimates for each survey month were subsequently generated by multiplying the mean number of animals per image by the total number of images required to cover the Survey Area.
47. Non-parametric bootstrap methods were used for variance estimation. A variability statistic was generated by re-sampling 999 times with replacement from the raw count data. The statistic was evaluated from each of these 999 bootstrap samples and upper and lower 95% confidence intervals of these 999 values were taken as the variability of the statistic over the population (Efron and Tibshirani 1993).

48. A measure of precision was calculated using a Poisson estimator, suitable for a pseudo-Poisson over-dispersed distribution. This produced a coefficient of variation (CV) based on the relationship of the standard error to the mean.
49. Species distributions maps were created for each species for each survey using QGIS.
50. **Table 11-4** and **Table 11-5** shows the raw numbers of marine mammals recorded during the digital aerial surveys in DBS East and West, respectively, from March 2021 to February 2022. These numbers are the raw count, therefore only present a relative abundance and not total abundances. The results indicate that harbour porpoise is present in the highest numbers, followed by dolphin or porpoise species which at worst case scenario can be considered as harbour porpoises. Unidentified seals are present in the high numbers, similar to dolphin/porpoise species followed by grey seals, white beaked dolphins, common dolphins, unidentified dolphin and with just one individual identified as a minke whale.

Table 11-4 APEM Surveys Species Counts for DBS East Array Area and 4km Buffer (March 2021 to February 2022)

Survey date	Grey seal	Seal species	Harbour porpoise	Dolphin / porpoise species	White-beaked dolphin	Dolphin species	Minke whale	Marine mammal species
Mar-21	0	3	1	4	0	1	0	0
Apr-21	0	0	8	9	0	0	0	0
May-21	0	2	14	4	0	0	0	0
Jun-21	2	3	39	2	0	0	0	0
Jul-21	2	4	39	0	0	0	0	1
Aug-21	3	0	21	1	0	0	0	1
Sep-21	1	1	64	2	0	0	1	0
Oct-21	1	1	12	0	0	0	0	0
Nov-21	0	1	18	0	0	0	0	0
Dec-21	0	6	12	9	8	0	0	2
Jan-22	1	2	27	4	0	0	0	0
Feb-22	1	2	21	0	0	0	0	0

Table 11-5 APEM Surveys Species Counts for DBS West Array Area and 4km Buffer (March 2021 to February 2022)

Survey date	Grey seal	Seal species	Harbour porpoise	Dolphin / porpoise species	White-beaked dolphin	Common dolphin	Dolphin species	Marine mammal species
Mar-21	0	0	0	5	0	0	1	0
Apr-21	1	1	8	7	4	0	1	1
May-21	0	9	48	5	0	2	0	2
Jun-21	6	10	97	7	8	0	0	3
Jul-21	8	5	52	1	0	0	0	1
Aug-21	4	0	22	1	0	0	0	0
Sep-21	0	1	29	4	0	0	0	1
Oct-21	1	1	8	0	0	0	0	0
Nov-21	0	0	13	0	0	0	0	4
Dec-21	3	3	8	4	0	0	0	0
Jan-22	3	3	29	0	0	0	0	0
Feb-22	5	1	30	0	0	0	0	4
Total	31	34	344	34	12	2	2	16

51. From the sightings numbers (as shown above) of each marine mammal species, or marine mammal species group, abundance and density estimates were calculated. Upper and lower confidence intervals (CI) as well as CV were also calculated for these density and abundance estimates. The density of animals at the sites (and hence the population size), the standard deviation, 95% CI and CV are then estimated using a non-parametric bootstrap method with replacement (Canty and Ripley 2010).

52. For species, such as marine mammals, that dive and therefore spend a considerable amount of time underwater, an availability bias, or correction factor, must be applied in order to account for those individuals that it is not possible to survey as they are underwater.
53. Without these availability bias, or correction factors, being corrected for, any abundance or density estimate would be relative only, rather than being an absolute estimate. For the site specific surveys, APEM have not calculated any correction factors for marine mammal data.
54. Species distribution maps have also been generated from the site-specific survey data at the Projects.

11.5.2 Surveys Within the Dogger Bank Area

55. HiDef Aerial Surveying Limited (HiDef) carried out digital aerial video surveys over the Round 3 Dogger Bank zone with additional effort focused on Tranche C, see **Plate 11-4**. The surveys were used to generate species density and distribution maps for the area with a 4km buffer. Surveys took place from January to December 2013. Transects were placed 7.5km apart in the main Dogger Bank zone, and 2.5km apart in Tranche C. Twelve surveys were undertaken between January 2013 to December 2013.
56. Harbour porpoises were the most numerous marine mammals recorded (n=2,702) with high numbers were recorded in March to September, with a peak in July (n=1,095). Grey seals were recorded in all surveys (n=92), peaking in January, March, and June 2013. White-beaked dolphins were recorded in most months (n=56), peaking in May and July 2013. Minke whale were recorded May to September (n=8) and the total for harbour seal was (n=12), seal species (n=59), unidentified dolphin (n=4), small cetacean/seal species (n=82) and unidentified cetacean (n=18).
57. Digital aerial video surveys over the Round 3 Dogger Bank zone were also carried out between April 2010 to May 2012. Again, harbour porpoise was the most numerous marine mammal recorded (N=5,728), recorded in every survey, peaks in the spring and summer. White beaked dolphin was recorded (n=104). White-beaked dolphin was recorded in May and June 2010, January, March to May, November and December 2011, March, and April 2012. Grey seal were more present in the summer months (N=46). Minke whale were recorded in spring and summer (n=18). Additional sightings that could not be attributed to a species small cetacean (n=6,034); dolphin species (n=89); seal species (n=218); small cetacean/seal species (n=210); large cetacean (m=9), unidentified cetacean (n=5) and cetacean/seal/shark (n=241).




Plate 11-4 Location of Aerial Survey Transects Within Dogger Bank and Tranche C.

58. The abundance of each species observed within the survey area was estimated separately using a design-based strip transect analysis with variance and confidence intervals derived through bootstrapping.
59. Gardline Environmental Ltd carried out boat based ornithology surveys for the Forewind Round 3 Dogger Bank zone (**Plate 11-5**).




Plate 11-5 Boat-based Survey Transect Lines in Dogger Bank (Gardline)

60. ESAS trained Seabird Surveyors undertook intensive boat-based surveys using the Camphuysen approach (Camphuysen *et al.* 2004) as recommended by COWRIE (Collaborative Offshore Wind Research into the Environment). The methods were based on distance sampling (Buckland *et al.* 2001; Stone *et al.* 1995), permitting accurate density estimation. There were 41 transect lines in total, with 4km spacing.

61. In addition to seabird's, incidental marine mammal sightings were also recorded. White-beaked dolphin were the most numerous marine mammal (n=811), peaking in April to June each year; followed by harbour porpoise (n=703), with no obvious peaks or trends; grey seal (n=244) and minke whale (n=199), sightings in spring and summer. Other species included unidentified dolphin (n=64), bottlenose dolphin (n=25), Atlantic white-sided dolphin (n=5); sela species (n=19); harbour seal (n=15); baleen whale species (n=7), humpback whale (n=4) and fin whale (n=2). The number of odontocetes recorded during the surveys are presented in **Plate 11-6**.




Plate 11-6 Incidental Odontocete Sightings During Ornithology Surveys for the Forewind Dogger Bank Offshore Wind Farm. January 2010 to June 2012

11.5.4 Other Available Sources

62. Other sources that have been used to inform the assessment are listed in **Table 11-6**.

Table 11-6 Other Available Data and Information Sources

Data Set	Spatial Coverage	Year	Notes
Small Cetaceans in the European Atlantic and North Sea (SCANS-III) data (Hammond <i>et al.</i> 2021)	North Sea and European Atlantic waters	Summer 2016	Provides information including abundance and density estimates of cetaceans in European Atlantic waters in summer 2016, including the offshore project areas.
Management Units (MUs) for cetaceans in UK waters (IAMMWG, 2022)	UK waters	2022	Provides information on MU for the offshore project areas.
Offshore Energy Strategic Environmental Assessment (OESEA) (including relevant appendices and technical reports) (OESEA 3 (DECC, 2016); OESEA 4 (Business, Energy and Industrial Strategy (BEIS), 2022a)).	UK waters	2016 2022	Provides information on marine mammals in UK waters.
The identification of discrete and persistent areas of relatively high harbour porpoise density in the wider UK marine area (Heinänen and Skov, 2015)	UK waters	1994-2011	Data was used to determine UK harbour porpoise Special Area of Conservation (SAC) sites.
Revised Phase III data analysis of Joint Cetacean Protocol (JCP) data resources (Paxton <i>et al.</i> 2016)	UK EEZ	1994-2011	Provides information on cetaceans in UK waters.
Seasonal habitat-based density models for a marine top predator, the harbour porpoise, in a dynamic environment (Gilles <i>et al.</i> 2016)	UK (SCANS II, Dogger Bank), Belgium, the Netherlands, Germany, and Denmark	2005-2013	Provides information for central and southern North Sea.

Data Set	Spatial Coverage	Year	Notes
Distribution and abundance maps for cetacean species around Europe (Waggitt <i>et al.</i> 2019).	North-east Atlantic	1980-2018	Provides information on cetacean species in North Sea and UK waters.
Distribution of Cetaceans, Seals, Turtles, Sharks and Ocean Sunfish recorded from Aerial Surveys 2001-2008 (The Wildfowl and Wetlands Trust (WWT), 2009)	UK areas of the North Sea	2001-2008	Provides information on species in the North Sea.
Sea Watch Foundation volunteer sightings off eastern England (Sea Watch Foundation, 2022)	East coast of England	2019-2020	Provides information on species sighted along east coast of England.
Habitat-based predictions of at-sea distribution for grey and harbour seals in the British Isles (Carter <i>et al.</i> 2020 and 2022)	North Sea	1991-2019	Provides information on abundance and absolute density estimates (i.e. number of seals) for seal species.
Seal telemetry data (e.g. Russell and McConnell, 2014; Russell, 2016a; Carter <i>et al.</i> 2020)	North Sea	1988-2010; 2015	Provides information on movements and distribution of seal species.
Special Committee on Seals (SCOS) annual reporting of scientific advice on matters related to the management of seal populations (SCOS, 2020; SCOS, 2021).	North Sea	2019 & 2020	Habitat-based predictions of at-sea distribution for grey and harbour seals in the British Isles.
Counts of grey seal in the Wadden Sea (Schop <i>et al.</i> 2022).	Wadden Sea	Winter 2021 to Spring 2022	Counts of grey seal during moult season.
Counts of harbour seal counts in the Wadden Sea (Galatius <i>et al.</i> 2022).	Wadden Sea	August 2022	Counts of harbour seal during pupping season.

11.6 Existing Environment

11.6.1 Harbour Porpoise

11.6.1.1 Desk-Based Review of Harbour Porpoise Presence

63. Heinänen and Skov (2015) identified one area of high harbour porpoise density in the summer period; from the western slopes of Dogger Bank south along a 30m depth contour towards an area off the Norfolk coast. High densities in winter were also identified in the southern North Sea, within an area between Flamborough Head and the outer Thames Estuary. High densities of harbour porpoise were predicted near both DBS East and DBS West, while high densities in summer were predicted to be further offshore.
64. The JCP Phase III Report (Paxton *et al.* 2016) identifies a similar distribution of high harbour porpoise density, with a relatively high density in the southern North Sea, with an estimated density of 8.1 to 18.3 individuals per km² in the vicinity of the Offshore Development Area (2.4% MU; Paxton *et al.* 2016).
65. Seasonal maps produced by Gilles *et al.* (2016) for harbour porpoise density across the central and south-eastern North Sea, indicated that in spring there were higher density areas in the southern and south-eastern part of the North Sea (with an estimated density of 0.53 individuals per km² in the vicinity of the Offshore Development Area). In summer, there was an apparent shift, compared to spring, toward offshore and western areas (with an estimated density of 0.59 individuals per km² in the vicinity of the Offshore Development Area). In autumn, there were higher densities compared to spring and lower densities compared to summer, (with an estimated density of 0.58 individuals per km² in the vicinity the Offshore Development Area) and in the winter, same density as spring (with an estimated density of 0.53 individuals per km² in the vicinity of the Offshore Development Area).
66. Distribution and abundance maps were developed by Waggitt *et al.* (2019) for cetacean species around Europe. For harbour porpoise, the distribution maps show a clear pattern of high harbour porpoise density in the southern North Sea, and the coasts of south-east England, for both January and July (**Plate 11-7**, Waggitt *et al.* 2019). Interrogation of this data², including all 10km 'grids' that overlap with the specified area, reveals an average annual density estimate of:

² Available from: [https:// doi.org/10.5061/dryad.mw6m905sz](https://doi.org/10.5061/dryad.mw6m905sz)

- 0.59 individuals per km² (average of all overlapping 10km 'grids') for the DBS East array area;
- 0.58 individuals per km² (average of all overlapping 10km 'grids') for the DBS West array area; 0.56 individuals per km² (average of all overlapping 10km 'grids') for the offshore export cable corridor; and
- 0.415 individuals per km² for the total Offshore Development Area.




Plate 11-7 Spatial Variation in Predicted Densities (Individuals per km of Harbour Porpoise in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.

11.6.1.2 Results from the Site-Specific Surveys for Harbour Porpoise

67. Data from the DBS array areas specific surveys were used to generate species distribution maps for both DBS array areas with a 4km buffer.
68. As noted above, harbour porpoise was the most commonly sighted marine mammal species during the surveys, with a total of 276 individuals recorded through the 12 survey dates at DBS East array area and 344 individuals recorded through the 12 survey dates at DBS West array area. Totalling at 620 individuals at DBS.
69. The distribution of harbour porpoise within DBS array areas varied, with individuals present across both DBS East array area and DBS West array area). There is no evident pattern of harbour porpoise distribution within the DBS array area, therefore no indication of a particular area of importance.

11.6.1.3 Results from Surveys in the Dogger Bank Area

70. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), a total of 2,702 harbour porpoises were recorded.
71. Throughout the Round 3 Dogger Bank Zone site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), harbour porpoises were recorded every month, with a total of 5,728 harbour porpoises. In addition, there were 306 small cetaceans recorded which most likely can be attributed to harbour porpoise. No species abundance or density estimates were undertaken.
72. During the ornithological surveys for the Forewind Dogger Bank Zone (monthly surveys January 2010 to June 2012 (excluding February 2012)), there were 344 incidental sightings of harbour porpoises with 703 individuals. No species abundance or density estimates were undertaken.

11.6.1.4 Site-Specific Density Estimates for Harbour Porpoise

73. Density estimates of animals/km² have been calculated from the raw data counts for (i) harbour porpoise and (ii) dolphin/porpoise species and are set out below. These abundance and densities are for the DBS East array area and the 4km buffer (**Table 11-7; Table 11-9**) and DBS West array area and the 4km buffer (**Table 11-8; Table 11-10**). No correction factor has been used to count for diving porpoises, therefore the abundance and density estimates are the absolute minimum.
74. The abundance estimates of harbour porpoise and dolphin/porpoise species are presented on a graph in **Plate 11-8** for DBS East array area and **Plate 11-9** for DBS West array area.

Table 11-7 Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km²) of Harbour Porpoises in: a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer

a) DBS East array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	1	9	1	26	1.00	0.01
Apr-21	8	69	26	129	0.35	0.07
May-21	14	121	52	199	0.27	0.13
Jun-21	39	338	205	486	0.16	0.37
Jul-21	39	334	189	506	0.16	0.20
Aug-21	21	181	69	302	0.22	0.60
Sep-21	64	549	214	1029	0.13	0.11
Oct-21	12	101	42	168	0.29	0.18
Nov-21	18	162	72	271	0.24	0.11
Dec-21	12	103	43	172	0.29	0.11
Jan-22	27	234	138	337	0.19	0.25
Feb-22	21	180	112	258	0.22	0.20
b) DBS East array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Mar-21	0	0	0	0	-	-
Apr-21	2	17	2	42	0.71	0.03
May-21	8	68	25	127	0.35	0.14
Jun-21	21	180	86	283	0.22	0.36
Jul-21	12	102	42	169	0.29	0.21
Aug-21	18	154	60	281	0.24	0.31
Sep-21	45	384	85	818	0.15	0.78
Oct-21	3	25	3	58	0.58	0.05
Nov-21	4	36	4	81	0.50	0.07
Dec-21	8	68	26	119	0.35	0.14
Jan-22	14	120	52	198	0.27	0.24
Feb-22	10	85	34	144	0.32	0.17

c) DBS East 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	1	9	1	26	1.00	0.02
Apr-21	6	52	9	104	0.41	0.12
May-21	6	53	9	107	0.41	0.12
Jun-21	18	158	79	264	0.24	0.37
Jul-21	27	235	122	383	0.19	0.55
Aug-21	3	26	3	70	0.28	0.06
Sep-21	19	164	52	311	0.23	0.38
Oct-21	9	77	26	145	0.33	0.18
Nov-21	14	127	45	227	0.27	0.30
Dec-21	4	35	4	78	0.50	0.08
Jan-22	13	113	52	183	0.28	0.26
Feb-22	11	96	43	157	0.3	0.22

Table 11-8 Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km²) of Harbour Porpoises in: a) DBS West Array Area plus 4km buffer; b) DBS West Array Area c) 4km buffer

a) DBS West array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	8	68	26	128	0.35	0.07
May-21	48	400	283	550	0.14	0.44
Jun-21	97	808	625	1017	0.10	0.88
Jul-21	52	436	285	596	0.14	0.48
Aug-21	22	184	84	301	0.21	0.20
Sep-21	29	238	107	402	0.19	0.26
Oct-21	8	68	17	143	0.35	0.07
Nov-21	13	110	51	177	0.28	0.12
Dec-21	8	67	25	126	0.35	0.07
Jan-22	29	244	143	344	0.19	0.27
Feb-22	30	253	160	346	0.18	0.28

b) DBS West array area						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	4	34	8	68	0.50	0.07
May-21	21	174	91	266	0.22	0.35
Jun-21	61	503	346	667	0.13	1.02
Jul-21	40	334	217	476	0.16	0.67
Aug-21	14	116	41	207	0.27	0.23
Sep-21	20	162	49	291	0.22	0.33
Oct-21	7	59	8	126	0.38	0.12
Nov-21	7	58	17	117	0.38	0.12
Dec-21	5	42	8	91	0.45	0.08
Jan-22	14	116	58	191	0.27	0.23
Feb-22	13	109	58	167	0.28	0.23
c) DBS West 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	4	34	4	86	0.50	0.08
May-21	27	226	134	343	0.19	0.54
Jun-21	36	304	203	431	0.17	0.72
Jul-21	12	101	34	186	0.29	0.24
Aug-21	8	68	8	144	0.35	0.16
Sep-21	9	75	17	167	0.33	0.18
Oct-21	1	9	1	26	1.00	0.02
Nov-21	6	51	17	102	0.41	0.12
Dec-21	3	25	3	59	0.58	0.06
Jan-22	15	128	60	214	0.26	0.30
Feb-22	17	145	77	222	0.24	0.34

Table 11-9 Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km²) of Unidentified Dolphin/Porpoise in a) DBS East Array Area plus 4km Buffer; b) DBS East Array Area c) 4km Buffer

a) DBS East array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	4	34	4	77	0.50	0.04
Apr-21	9	77	26	137	0.33	0.08
May-21	4	35	9	69	0.50	0.04
Jun-21	2	17	0	43	0.71	0.02
Aug-21	1	9	1	26	1.00	0.01
Sep-21	2	17	2	43	0.71	0.02
Dec-21	9	77	26	146	0.33	0.08
Jan-22	4	35	4	104	0.50	0.04
b) DBS East array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	42	0.74	0.03
Apr-21	5	42	8	93	0.45	0.09
May-21	0	0	0	0	-	-
Jun-21	0	0	0	0	-	-
Aug-21	0	0	0	0	-	-
Sep-21	1	9	1	26	1.00	0.02
Dec-21	3	26	3	68	0.58	0.05
Jan-22	1	9	1	26	1.00	0.02
c) DBS East 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	52	0.71	0.04
Apr-21	4	35	9	69	0.50	0.08
May-21	4	36	9	71	0.50	0.08
Jun-21	2	18	2	44	0.71	0.04
Aug-21	1	9	1	26	1.00	0.02
Sep-21	1	9	1	26	1.00	0.02
Dec-21	6	52	9	104	0.41	0.12
Jan-22	3	26	3	79	0.58	0.06

Table 11-10 Raw Counts, Abundance and Density Estimates (no. Estimated Individuals per km²) of Unidentified Dolphin/Porpoise in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer

a) DBS West array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	5	42	8	84	0.45	0.05
Apr-21	7	59	17	119	0.38	0.06
May-21	5	42	8	83	0.45	0.05
Jun-21	7	58	17	117	0.38	0.06
Jul-21	1	8	1	25	1.00	0.01
Aug-21	1	8	1	25	1.00	0.01
Sep-21	4	33	4	74	0.50	0.04
Dec-21	4	34	4	67	0.50	0.04
b) DBS West array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	42	0.71	0.03
Apr-21	5	42	8	84	0.45	0.08
May-21	3	25	3	58	0.58	0.05
Jun-21	7	58	16	107	0.38	0.12
Jul-21	1	8	1	25	1.00	0.02
Aug-21	1	8	1	25	1.00	0.02
Sep-21	4	32	4	73	0.50	0.06
Dec-21	3	25	3	58	0.58	0.05
c) DBS West 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	3	25	3	59	0.58	0.06
Apr-21	2	17	2	43	0.71	0.04
May-21	2	17	2	42	0.71	0.04
Jun-21	0	0	0	0	-	-
Jul-21	0	0	0	0	-	-
Aug-21	0	0	0	0	-	-
Sep-21	0	0	0	0	-	-
Dec-21	1	8	1	25	1.00	0.02

Dogger Bank South Offshore Wind Farms




Plate 11-8 Estimated Abundance of Harbour Porpoise Within DBS East Array Area.




Plate 11-9 Estimated Abundance of Harbour Porpoise Within DBS West Array Area

75. The results from the site-specific surveys provide seasonal absolute density estimates for harbour porpoise which are presented in **Table 11-11**.

Table 11-11 Seasonal Density Estimates for Harbour Porpoise from APEM Ltd Survey

Season	DBS East array area absolute density estimates	DBS West array area absolute density estimates
Summer average	0.275	0.427
Winter average	0.17	0.058
Yearly average	0.223	0.243

11.6.1.5 Review of Abundance and Density Estimates for Harbour Porpoise

76. A series of large-scale surveys for cetaceans in European Atlantic waters was initiated in summer 1994, in the North Sea and adjacent waters (SCANS, 1995; Hammond *et al.* 2002) and continued in summer 2005 in all shelf waters (SCANS-II 2008; Hammond *et al.* 2013). Despite no overall change in population size between the SCANS-I and SCANS-II surveys, large scale changes in the distribution of harbour porpoise were observed between 1994 and 2005, with the main concentration shifting from north-eastern UK and Denmark to the southern North Sea. Such large-scale changes in the distribution of harbour porpoise are likely the result of changes to the availability of principal prey within the North Sea (SCANS-II, 2008).
77. Calculating Waggitt *et al.* 2019 densities with all 10km 'grids' that overlap with the specified area, reveals an average annual density estimate of 0.59 individuals per km² for the DBS East array area; 0.58 individuals per km² for the DBS West array area; and 0.56 individuals per km² (average of all overlapping 10km 'grids') for the Offshore Development Area.
78. Results from the SCANS-III survey (the most recent available; undertaken in summer 2016; Hammond *et al.* 2021) also indicate that the occurrence of harbour porpoise is greater in the central and southern areas of the North Sea compared to the northern North Sea.
79. Within the impact assessments for harbour porpoise, in addition to the site specific density estimates for harbour porpoise, density estimates from the SCANS-III surveys (Hammond *et al.* 2021) will also be used to provide context for the wider area. The DBS sites are both in SCANS-III survey block O (**Plate 11-10; Plate 11-11**):
 - Abundance = 53,485 harbour porpoise (CV = 0.209; 95% CI = 37,413-81,695)
 - Density = 0.888 harbour porpoise/km² (CV=0.209)




Plate 11-10 Area Covered by the SCANS-III Survey, and the Locations of the SCANS-III Survey Blocks (The Projects Are Within Survey Block O) (Hammond et al. 2021)




Plate 11-11 Estimated Harbour Porpoise Density in Each SCANS-III Survey Block (Hammond *et al.* 2021)

80. Harbour porpoise within the eastern North Atlantic is generally considered to be part of a continuous biological population that extends from the French coastline of the Bay of Biscay to northern Norway and Iceland (Tolley and Rosel 2006; Fontaine *et al.* 2007, 2014; IAMMWG 2022). However, for conservation and management purposes, it is necessary to consider this population within smaller MUs. MUs provide an indication of the spatial scales at which effects of plans and projects alone, and in-combination, need to be assessed for the key cetacean species in UK waters, with consistency across the UK (IAMMWG, 2022).

81. IAMMWG defined three MUs for harbour porpoise: North Sea (NS); West Scotland (WS); and the Celtic and Irish Sea (CIS). DBS is located in the NS MU (**Plate 11-12**).
82. The estimate of harbour porpoise abundance in the North Sea MU is 346,601 (CV = 0.09; 95%; CI = 289,498 – 419,967 IAMMWG 2022). This is the reference population for harbour porpoise, of which any potential impacts will be assessed against.




Plate 11-12 Harbour porpoise Management Units (MUs), noting that this species is largely confined to the continental shelf (i.e. waters <200m depth) (IAMMWG, 2022).

11.6.1.6 Diet of Harbour Porpoise

83. The distribution and occurrence of harbour porpoise, as well as other marine mammal species is most likely to be related the availability and distribution of their prey species. For example, sandeels *Ammodytidae* species, which are known prey for harbour porpoise, exhibit a strong association with key surface sediments (Gilles *et al.* 2016; Clarke *et al.* 1998).

84. Harbour porpoise are generalist feeders, and their diet reflects available prey in an area. Therefore, their diet varies geographically, seasonally and annually, reflecting changes in available food resources and differences in diet between sexes or age classes may also exist. The diet of the harbour porpoise consists of a wide variety of fish, including pelagic schooling fish, as well as demersal and benthic species, especially Gadoids, Clupeids and sandeels (Berrow and Rogan 1995; Kastelein *et al.* 1997; Börjesson *et al.* 2003; Santos and Pierce 2003; Santos *et al.* 2004).
85. Harbour porpoise tend to concentrate their movements in small focal regions (Johnston *et al.* 2005), which often approximate to particular topographic and oceanographic features and are associated with prey aggregations (Raum-Suryan and Harvey 1998; Johnston *et al.* 2005; Keiper *et al.* 2005; Tynan *et al.* 2005). Consequently, habitat use is highly correlated with prey density rather than any particular habitat type.
86. Harbour porpoise have relatively high daily energy demands and need to capture enough prey to meet its daily energy requirements. It has been estimated that, depending on the conditions, harbour porpoise can rely on stored energy (primarily blubber) for three to five days, depending on body condition (Kastelein *et al.* 1997).

11.6.2 Bottlenose Dolphin

11.6.2.1 Desk-Based Review of Bottlenose Dolphin Presence

87. Throughout its range, the bottlenose dolphin occurs in a diverse range of habitats, from shallow estuaries and bays, coastal waters, continental shelf edge and deep open offshore ocean waters. However, it is primarily an inshore species, with most sightings within 10km of land, but they can also occur offshore, often in association with other cetaceans³.
88. In coastal waters, bottlenose dolphin is often associated with river estuaries, headlands or sandbanks, where there is uneven bottom relief and/or strong tidal currents (e.g. Lewis and Evans 1993; Wilson *et al.* 1997; Liret *et al.* 1998; Liret, 2001; Ingram and Rogan 2002; Reid *et al.* 2003).

³ <https://sac.jncc.gov.uk/species/S1349/>

89. A resident population of bottlenose dolphin is present in the Moray Firth, with an estimated 224 individuals (95% CI 213 – 234; IAMMWG, 2022). Historically, very few sightings of bottlenose dolphin were recorded further south on the east coast of the UK. In recent years an increase in bottlenose dolphins along the coastline of north-east England have been reported (Aynsley, 2017; Hacket, 2022). They have been recorded approximately 300 miles outside of what would be considered their ‘normal’ home range (Cheney *et al.* 2018), with one individual from the Moray Firth population being recorded as far south and east as The Netherlands (Aynsley, 2017). Whilst bottlenose dolphin presence has been increasing in north-east England in recent years, they appear to be a coastal population at present (Hacket, 2022).
90. Bottlenose dolphin sightings were made year-round along the north-east England coast (between 2013 and 2016; Aynsley 2017), suggesting that there is no seasonal pattern to the increase in recent sightings numbers. A total of 48 of the individuals sighted within this period on the north-east coast were attributed to being part of the Moray Firth population using photo-identification. A total of 11 calves were also sighted, indicating that bottlenose dolphin in this area of coastline is reproductively active.
91. The results of the JCP Phase III Report (Paxton *et al.* 2016) identified that for bottlenose dolphin, densities are low across much of UK waters, with higher densities off the west coast of Wales, and within the Moray Firth. The density of bottlenose dolphin within the southern North Sea for DBS is low, with 0.5 to 1.7 individuals per km² (2.6% MU) (Paxton *et al.* 2016).
92. The SCANS-III survey shows a similar distribution pattern, with no bottlenose dolphin identified within the southern North Sea survey block L or the more northerly block O, and higher densities with block R, for the east coast of Scotland (**Plate 11-10; Plate 11-13**, Hammond *et al.* 2021).




Plate 11-13 Estimated Bottlenose Dolphin Density in Each SCANS-III Survey Block (Hammond *et al.* 2021)

93. For bottlenose dolphin, the distribution maps (developed by Waggitt *et al.* 2019) show a clear pattern of higher density to the western coastal areas of the UK, extending south to the Bay of Biscay (**Plate 11-14**, Waggitt *et al.* 2019). Densities of bottlenose dolphin in the North Sea are very low in comparison. Interrogation of this data⁴, including all 10km 'grids' that overlap with the specified area, reveals an average annual density estimate of:

- 0.00014 individuals per km² for the DBS East array area;
- 0.00025 individuals per km² for the DBS West array area; and
- 0.0013 individuals per km² for the offshore export cable corridor; and
- 0.0014 individuals per km² for the total Offshore Development Area.




Plate 11-14 Spatial Variation in Predicted Densities (Individuals per km of Bottlenose Dolphin in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.

⁴ Available from: [https:// doi.org/10.5061/dryad.mw6m905sz](https://doi.org/10.5061/dryad.mw6m905sz)

11.6.2.2 Results from the Site-Specific Surveys for Bottlenose Dolphin

94. During the site specific digital aerial surveys of both DBS East array area and DBS West array area, undertaken from March 2021 to February 2022, no bottlenose dolphin were recorded. However, one sighting was recorded as unidentified dolphin, which could have been attributed to bottlenose dolphin.

11.6.2.3 Results from Surveys in the Dogger Bank Area

95. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), no bottlenose dolphins were recorded. However, during the 12 surveys, 190 sightings were recorded with 'no species ID' some of which were further recorded as a cetacean/small cetacean which could possibly be bottlenose dolphins. No abundance or density estimated were calculated for unknown species.
96. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), no bottlenose dolphins were recorded. In addition, there were 306 small cetaceans recorded which possibly can be attributed to bottlenose dolphins, along with 89 unidentified delphinid species. No species abundance or density estimates were undertaken.
97. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there was one sighting of bottlenose dolphins with 25 individuals. There were also 24 incidental sightings of unidentified delphinid species, and one acoustic detection, resulting in 64 individuals which could possibly be attributed to bottlenose dolphins. No species abundance or density estimates were undertaken.

11.6.2.4 Site-Specific Density Estimates for Bottlenose dolphin

98. Based on the DBS site specific surveys, density estimates of animals/km² have been calculated from the raw data counts for unidentified dolphin species as there is a possibility that these can be bottlenose dolphins. In total, one unidentified dolphin was recorded during the April 2021 survey, resulting in an abundance estimate of nine and density of 0.01 marine mammal/km².

11.6.2.5 Review of Abundance and Density Estimates for Bottlenose Dolphins

99. As sightings of bottlenose dolphin have been increasingly reported along the north-east coast of England, they have also been included in the assessment. For the entire SCANS-III survey area, bottlenose dolphin abundance in the summer of 2016 was estimated to be 19,201, with an overall estimated density of 0.00159/km² (CV = 0.242; 95% CI = 11,404-29,670; Hammond *et al.* 2021).

100. There is currently no density estimate for bottlenose dolphin in and around the Projects based on the SCANS-III density estimates. The adjacent survey block R to the North which covers the Moray Firth area is thought to be the same bottlenose dolphin population as has recently been recorded off the east coast of England.
- Abundance = 1,924 bottlenose dolphin (CV=0.861; 95% CI=0-5,048)
 - Density = 0.0298 bottlenose dolphin/km² (CV=0.861)
101. As for the density estimate, the GNS MU, which the Projects are located within, has a population estimate for the bottlenose dolphin of 2,022 (95% CI = 548 – 7,453; IAMMWG 2022). The results from Waggitt *et al.* (2019) seasonal absolute density estimates for bottlenose dolphin are presented in **Table 11-12**.

Table 11-12 Seasonal Density Estimates for Bottlenose Dolphin from Waggitt *et al.* (2019) Densities

Season	DBS East array area absolute density estimates	DBS West array area absolute density estimates	Offshore Development Area
Summer average	0.00014	0.00025	0.0013
Winter average	0.00009	0.00020	0.0011
Yearly average	0.00011	0.00023	0.0012

102. Therefore, within the impact assessments for bottlenose dolphin, the worst-case density estimates relative to the Projects from the Waggitt *et al.* 2019 data will be used in the assessment:
- 0.00014 individuals per km² for the DBS East array area;
 - 0.00025 individuals per km² for the DBS West array area;
 - 0.0013 individuals per km² for the offshore export cable corridor; and
 - 0.0014 individuals per km² for the total Offshore Development Area.




Plate 11-15 Bottlenose Dolphin MU (IAMMWG, 2022).

11.6.2.6 Diet of Bottlenose Dolphin

103. Bottlenose dolphins are opportunistic feeders and take a wide variety of fish and invertebrate species. Benthic and pelagic fish (both solitary and schooling species), including haddock *Melanogrammus aeglefinus*, saithe *Pollachius virens*, pollock *Pollachius pollachius*, cod *Gadus morhua*, whiting *Merlangius merlangus*, hake *Merluccius merluccius*, blue whiting *Micromesistius poutassou*, bass *Dicentrarchus labrax*, mullet *Mugilidae*, mackerel *Scombridae*, salmon *Salmo salar*, sea trout *Salmo trutta trutta*, flounder *Platichthys flesus*, sprat *Sprattus sprattus* and sandeels, as well as octopus and other cephalopods have all been recorded in the diet of bottlenose dolphin (Santos *et al.* 2001; Santos *et al.* 2004; Reid *et al.* 2003).

104. Diet analysis suggests that bottlenose dolphin are selective opportunists and although they may have preference for a type of prey, their diet seems to be determined largely by prey availability. Research in Australia has shown that when presented with a choice, they will preferentially feed on certain types of prey, particularly those with a high fat content (Corkeron *et al.* 1990).
105. Analysis of the stomach contents of ten bottlenose dolphin in Scottish waters, from 1990 to 1999, reveals that the main prey are cod (29.6% by weight), saithe (23.6% by weight), and whiting (23.4% by weight), although other species including salmon (5.8% by weight), haddock (5.4% by weight) and cephalopods (2.5% by weight) were also identified in lower number (Santos *et al.* 2001).

11.6.3 Common Dolphin

11.6.3.1 Desk-Based Review of Common Dolphin Presence

106. The short-beaked common dolphin is a widespread species of common dolphin occurring in warm-temperate oceans. Being one of the most abundant species of cetacean, the common dolphin has a global population estimated to be around six million (Hammond *et al.* 2008).
107. Throughout its range, the common dolphin is primarily distributed in the Celtic Sea and Western Approaches to the Channel, and off southern and western Ireland (BEIS 2022b; Hammond *et al.* 2021; Waggitt *et al.* 2019).
108. There is very little literature on common dolphins in the North Sea, however it is documented that they have a seasonal occurrence in the North Sea in the summer months (Waggitt *et al.* 2019).
109. Results from ORCA (*Organisation cetacea*) surveys carried out yearly have recorded 20 sightings of common dolphins with 52 individuals from 2006 to 2017 in the summer months in the North Sea (ORCA 2023).
110. The results of the JCP Phase III Report (Paxton *et al.* 2016) identified that for common dolphin, densities are high across much of UK waters, with 102,656 (0.29) MU in CGNS (**Plate 11-16**), but low in DBS <0.01 individuals per km² (0.000009% MU) (Paxton *et al.* 2016).
111. No common dolphins were recorded in survey block O (in which the Projects are located) or any of the blocks in the North Sea during the SCANS-III survey (**Plate 11-16**, Hammond *et al.* 2021).




Plate 11-16 Estimated Common Dolphin Density in Each SCANS-III Survey Block (Hammond *et al.* 2021)

112. For common dolphin, the distribution maps (developed by Waggitt *et al.* 2019) show a clear pattern of higher density to the western coastal areas of the UK, extending south to the Bay of Biscay (**Plate 11-17**; Waggitt *et al.* 2019). Densities of common dolphin in the North Sea are very low in comparison, with the peak recorded in the summer months. The distribution maps also indicate a 'corridor' of increased common dolphin density travelling from west of Scotland, southwards around the west coast of the Northern Ireland and the Republic of Ireland, and through the centre of the Bay of Biscay, with little occurrence in the North Sea.




Plate 11-17 Spatial Variation in Predicted Densities Individuals per km of Common Dolphin in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.

11.6.3.2 Results from the Site-Specific Surveys for Common Dolphin

113. During the site specific digital aerial surveys of both DBS East array area and DBS West array area, undertaken from March 2021 to February 2022, two common dolphin were recorded (**Table 11-13**).
114. In addition to the common dolphins; there was one sighting recorded as unidentified dolphin, which could have been attributed to common dolphin (**Table 11-9; Table 11-10**).

11.6.3.3 Results from Surveys in the Dogger Bank Area

115. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), no common dolphins were recorded. However, during the 12 surveys, 190 sightings were recorded with 'no species ID' some of which were further recorded as a cetacean/small cetacean which could possibly be attributed to common dolphins. No abundance or density estimated were calculated for unknown species.
116. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), no common dolphins were recorded. In addition, there were 306 small cetaceans recorded which possibly can be attributed to bottlenose dolphins, along with 89 unidentified delphinid species. No species abundance or density estimates were undertaken.
117. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there was one sighting of bottlenose dolphins with 25 individuals. There were also 24 incidental sightings of unidentified delphinid species, and one acoustic detection, resulting in 64 individuals which could possibly be attributed to common dolphin. No species abundance or density estimates were undertaken.

11.6.3.4 Site-Specific Density Estimates for Common Dolphin

118. **Table 11-13** presents the abundance and density estimates of common dolphins from the DBS site-specific surveys (APEM, 2022). Common dolphin was recorded in the DBS West array area, with one occurrence in March, with two individuals. Therefore the density estimate represent March 2021 and is zero for all other months of the first years survey.

Table 11-13 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km²) of Common Dolphins: a) DBS West Array Area Plus 4km; b) DBS West Array Area c) 4km Buffer

a) DBS West array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	42	0.71	0.02
b) DBS West array area						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	1	8	1	25	1	0.02

c) DBS West 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	1	8	1	25	1	0.02

11.6.3.5 Review of Abundance and Density Estimates for Common Dolphin

119. For both SANS-II and SCANS-III surveys, there are no abundance or density estimates for common dolphin in the North Sea (Hammond *et al.* 2021).
120. Calculating Waggitt *et al.* 2019 densities including all 10km 'grids' that overlap with the DBS areas, reveals an average annual density estimate of 0.037 individuals per km² for the DBS East array area; 0.037 individuals per km² for the DBS West array area; and 0.073 individuals per km² for the offshore export cable corridor.
121. For the site specific surveys (APEM Ltd), the abundance estimate for common dolphin is eight and the density is 0.02km².

Table 11-14 Seasonal Density Estimates for Common Dolphin from Waggitt *et al.* 2019 Densities

Season	DBS East array area absolute density estimates	DBS West array area absolute density estimates	Offshore Development Area
Summer average	0.037	0.037	0.073
Winter average	0.021	0.021	0.041
Yearly average	0.029	0.029	0.057

122. Therefore, within the impact assessments for common dolphin, density estimates from Waggitt *et al.* 2019 data will be used:
- 0.037 individuals per km² for the DBS East array area;
 - 0.037 individuals per km² for the DBS West array area;
 - 0.073 individuals per km² for the offshore export cable corridor and
 - 0.038 individuals per km² for the total Offshore Development Area.
123. For the density estimate, the CGNS MU (**Plate 11-18**) which the Projects are located in, has a population estimate for common dolphin of 102,656 (95% CI = 58,932 – 178,822; IAMMWG, 2022).




Plate 11-18 Common Dolphin, White-beaked Dolphin and Minke Whale MU (IAMMWG, 2022).

11.6.3.6 Diet of Common Dolphin

124. Common dolphin are cooperative feeders, working within a pod to capture prey. They have a varied diet of fish including haddock, mackerel *Scomber scombrus*, Atlantic horse mackerel *Trachurus trachurus*, blue whiting, anchovy *Engraulida spp.*, and sardine *Sardina pilchardus* (Couperus 1997; Silva 1999; Meynier 2004) which are also exploited by fisheries. Other prey items recorded in common dolphins include cephalopods and crustacean (Brophy *et al.* 2009).
125. Analysis of 63 common dolphin stomach contents from the Bay of Biscay found that the diet was dominated by fish with mackerel being the preferred fish and cephalopods were recorded as a prey of secondary importance (Pusineri *et al.* 2007).

11.6.4 White-Beaked Dolphin

11.6.4.1 Desk-Based Review of White-Beaked Dolphin Presence

126. White-beaked dolphin are widely distributed within the central North Sea, however, very few sightings are recorded along the east coast of England or south of the Humber Estuary, with a small number of sightings in offshore waters within the shallow waters near the North Norfolk Sandbanks and Dogger Bank areas (Gilles *et al.* 2012; DECC 2016). The occurrence of white-beaked dolphin in the southern North Sea is relatively low (Reid *et al.* 2003; Hammond *et al.* 2013; 2021).
127. A review of the strandings data of white-beaked dolphin in the North Sea were collated and assessed by ASCOBANS (IJseldijk *et al.* 2018) in order to determine temporal and spatial trends in the distributions of white-beaked dolphin in the south-western North Sea. Strandings data used within the review were from Belgium, Germany, the Netherlands and the UK, from 1991 to 2017. This review indicates that there has been a reduction in the abundance of white-beaked dolphin in the south-east coasts of the UK, with an increase in the north-east area (IJseldijk *et al.* 2018).
128. Data on the distribution of marine mammals in UK areas of the North Sea have been collected opportunistically during aerial surveys for birds conducted by WWT Consulting from 2001-2008 (WWT 2009). A number of unknown dolphin species were recorded, with local clusters present north-east off Flamborough Head. White-beaked dolphin were also recorded in small numbers in the north-east, again off Flamborough Head (WWT 2009).
129. The results of the JCP Phase III Report (Paxton *et al.* 2016) identified that for white-beaked dolphin, densities are low across much of UK waters, with higher densities shown to be in the Hebrides and the northern North Sea. The density of white-beaked dolphin within the southern North Sea in the vicinity of the Offshore Development Area is 0.1 to 2.7 individuals per km² (1.3% MU; Paxton *et al.* 2016).
130. The SCANS-III survey shows a similar distribution pattern, with no white-beaked dolphin identified within the southern North Sea survey block L, and low but increasing densities with the more northerly North Sea survey blocks (blocks O and R) (Hammond *et al.* 2021).
131. For white-beaked dolphin, the distribution maps (developed by Waggitt *et al.* 2019) show a clear pattern of higher density in the northern North Sea, and around the coasts of Scotland, with decreasing densities southwards of Scotland along the east coast of England. There is also a clear seasonal difference in the densities of white-beaked dolphin, with higher densities in July, particularly to the north of their range (**Plate 11-19**; Waggitt *et al.* 2019).




Plate 11-19 Spatial Variation in Predicted Densities Individuals per km of White-beaked Dolphin in January and July in the NE Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.

11.6.4.2 Results from the Site-Specific Surveys for White-Beaked Dolphin

132. During the site-specific aerial surveys of both DBS East and DBS West array areas, undertaken from March 2021 to February 2022, eight white-beaked dolphins were recorded in DBS East array area during one survey and 12 individuals in DBS West array area during two surveys. However, several sightings were recorded as unidentified dolphin, dolphin/porpoise species and marine mammal species, some of which could have been white-beaked dolphin (**Table 11-4, Table 11-5**).

11.6.4.3 Results from Surveys in the Dogger Bank Area

133. Data from the Dogger bank Tranche C site specific surveys with a 4km buffer (12 surveys undertaken between January 2013 to December 2013); 56 individuals were recorded through the 12 survey dates, peaking May to July.

134. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), a total of 104 white-beaked dolphins were recorded, peaking in the spring. In addition, there were 306 small cetaceans recorded which possibly can be attributed to white-beaked dolphins, along with 89 unidentified delphinid species. No species abundance or density estimates were undertaken.
135. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there were 138 incidental sightings of white-beaked dolphins, with 811 individuals. There was also a further 24 sightings of unidentified delphinid species, which could be attributed to white-beaked dolphins. No species abundance or density estimates were undertaken.
136. Abundance and density estimate of animals/km² have been calculated from the raw data counts for (i) white-beaked dolphin and (ii) dolphin species which can be attributed to white-beaked dolphin. White-beaked dolphins were recorded in DBS East during the December 2021 survey, with eight individuals resulting in 0.07 dolphins/km². In DBS West, white-beaked dolphins were recorded in April and June 2012, resulting in a density estimate of 0.07 dolphins/km².

11.6.4.4 Site-Specific Density Estimates for White-Beaked Dolphin

137. Abundance and density estimate of animals/km² have been calculated from the raw data counts for (i) white-beaked dolphin and (ii) dolphin species which can be attributed to white-beaked dolphin. White-beaked dolphins were recorded in DBS East during the December 2021 survey, with eight individuals resulting in 0.07 dolphins/km². In DBS West, white-beaked dolphins were recorded in April and June 2012, resulting in a density estimate of 0.07 dolphins/km².

11.6.4.5 Review of Abundance and Density Estimates for White-Beaked Dolphin

138. For the entire SCANS-III survey area, white-beaked dolphin abundance in the summer of 2016 was estimated to be 36,287 with an overall estimated density of 0.030/km² (CV = 0.29; 95% CI = 18,694-61,869; Hammond *et al.* 2017). DBS is located in SCANS-III survey block O (Hammond *et al.* 2021):
 - Abundance = 143 white-beaked dolphin (CV=0.97; 95% CI= 0-490)
 - Density = 0.002 white-beaked dolphin/km² (CV=0.97)
139. Scientific evidence supports the assumption that white-beaked dolphin from around the British Isles and North Sea represent one population, with movement between Scottish waters and the Danish North Sea and Skagerrak (Banguera-Hinestroza *et al.* 2010; IAMMWG 2022).

140. However, it is worth noting that this species usually occurs on the continental shelf (Reid *et al.* 2003; IAMMWG 2022). The UK EEZ white-beaked dolphin abundance is 34,025 (CV = 0.28; 95% CI = 20,026 – 57,807), which are derived from the SCANS-III (Hammond *et al.* 2017) and ObSERVE data (Rogan *et al.* 2018). The reference population for white-beaked dolphin in the CGNS MU is 43,951 animals (CV=0.22; 95% CI= 28,439 – 67,924; IAMMWG 2022). The results from Waggitt *et al.* 2019 seasonal absolute density estimates for white-beaked dolphin are presented in **Table 11-15**.

Table 11-15 Seasonal Density Estimates for White-beaked Dolphin from Waggitt *et al.* 2019 Densities.

Season	DBS East array area absolute density estimates	DBS West array area absolute density estimates	Offshore Development Area
Summer average	0.0055	0.012	0.046
Winter average	0.0048	0.009	0.035
Yearly average	0.0052	0.011	0.041

141. Within the impact assessments for white-beaked dolphin, the worse-case density estimates for the offshore sites will be used. Therefore, within the impact assessments the density estimates from Waggitt *et al.* 2019 data will be used:
- 0.0055 individuals per km² for the DBS East array area;
 - 0.012 individuals per km² for the DBS West array area;
 - 0.046 individuals per km² for the offshore export cable corridor; and
 - 0.017 individuals per km² for the total Offshore Development Area.

11.6.4.6 Diet of White-Beaked Dolphin

142. Analysis of the stomach contents of white-beaked dolphin have shown that the species feed on a wide range of fish and squid species, including cod, whiting, and hake (Kinze *et al.* 1997; Reeves *et al.* 1999). White-beaked dolphin have also been observed to associate with herring *Clupea harengus* (Harmer, 1927; Fraser, 1946; Evans, 1980) and mackerel (Evans *et al.* 1987) shoals, and anecdotal evidence from fisherman in Scotland suggests that individuals seen inshore may coincide with mackerel appearing in the same areas (Canning *et al.* 2008).

143. Dietary analysis for 22 white-beaked dolphin stranded around the UK coast between 1992 and 2003 (Canning *et al.* 2008) found that while a wide variety of prey species were identified, the majority of prey were from a much smaller number of species. Haddock and whiting were the most predominantly found, representing 43% and 24% respectively of the total reconstructed weight, with cod representing a further 11% of the total reconstructed weight.

11.6.5 Minke Whale

11.6.5.1 Desk-Based Review of Minke Whale Presence

144. Minke whales are widely distributed along the Atlantic seaboard of Britain and Ireland and throughout the North Sea. The JNCC Cetacean Atlas (Reid *et al.* 2003), indicates that minke whale occur regularly in the North Sea to the north of Humberside, but are comparatively scarce in the southern North Sea. Animals are present throughout the year, but most sightings are between May and September (Reid *et al.* 2003). DECC (2016) support this, stating that sightings rarely extend past Dogger Bank, but that occasional sightings of minke whale are made as far south as Flamborough Head and the north Humberside coastlines between July and October (DECC 2016).
145. Higher densities of minke whale have been recorded along the margins of Dogger Bank and adjacent areas in spring and summer (de Boer 2010; Gilles *et al.* 2012; Hammond *et al.* 2013). Few sightings of minke whale have been made further south of these areas and it is thought that they probably enter the North Sea from the north (DECC 2016). Minke whales appear to move into the North Sea at the beginning of May and are present throughout the summer until October (Northridge *et al.* 1995).
146. The JCP Phase III Report (Paxton *et al.* 2016) identified a total of 1,860 minke whale sightings within the UK offshore area. The density of minke whale was predicted to be highest along the northern coast of the UK, from Yorkshire north to the Kintyre Peninsula. The resultant density maps produced in the JCP Phase III Report (Paxton *et al.* 2016) show a minke whale density of less than 0.04 per km² for the southern North Sea (97.5% CI 0-0.02 – 0.08 per km²) below the Humber Estuary and Flamborough Head. The JCP report estimated 0.8 to 4.9 individuals per km² in the offshore development area (1.3% MU; Paxton *et al.* 2016).

147. For minke whale, the distribution maps (developed by Waggitt *et al.* 2019) show a clear pattern of higher density in the northern North Sea, and around the coasts of Scotland, Ireland and within the CIS, with decreasing densities southwards of Scotland along the east coast of England. There is a clear seasonal difference in the densities of minke whale, with higher densities in July, which is particularly evident in the north of their range (**Plate 11-20**; Waggitt *et al.* 2019).
148. In addition, the distribution maps indicate a ‘corridor’ of increased minke whale density travelling from north of Orkney, around the north and west coasts of the UK to Northern Ireland. DBS array areas are located to the very southern end of the area with relatively higher densities, and there appears to be a slight difference in their seasonal distributions with higher densities in this area during the summer months. Interrogation of this data⁵, including all 10km ‘grids’ that overlap with the specified area, reveals an average annual density estimate of:
- 0.0025 individuals per km² (average of all overlapping 10km ‘grids’) for the DBS East array area;
 - 0.0034 individuals per km² (average of all overlapping 10km ‘grids’) for the DBS West array area;
 - 0.0073 individuals per km² (average of all overlapping 10km ‘grids’) for the offshore export cable corridor; and
 - 0.0075 individuals per km² for the total Offshore Development Area.

⁵ Available from: [https:// doi.org/10.5061/dryad.mw6m905sz](https://doi.org/10.5061/dryad.mw6m905sz)




Plate 11-20 Spatial Variation in Predicted Densities (Individuals per km of Minke Whale in January and July in the North-East Atlantic). Values are Provided at 10km Resolution. Source: Waggitt et al. 2019.

11.6.5.2 Results from the Site-Specific Surveys for Minke Whale

149. During the DBS array areas site specific digital aerial surveys (12 surveys undertaken between March 2021 and February 2022), a single minke whale was positively identified in September inside DBS East site.

11.6.5.3 Results from Surveys in the Dogger Bank Area

150. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), eight minke whale were recorded.
151. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), a total of 18 minke whales were recorded, peaking in June 2010, and 2011. In addition, there were nine large cetaceans recorded which possibly can be attributed to minke whales. No species abundance or density estimates were undertaken.

152. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there were 183 incidental sightings of minke whales, with 199 individuals. There was also a further six sightings of solitary unidentified baleen whales, which most probably can be attributed to minke whales.

11.6.5.4 Site-Specific Density Estimates for Minke Whale

153. During the DBS array areas site specific digital aerial surveys (12 surveys undertaken between March 2021 and February 2022), the individual minke recorded, resulted in a relative density estimate of 0.01 individuals per km². This is the same density estimate as for the SCANS-III survey (**Section 11.6.5.2**). The minke whale was recorded in the DBS East array area.
154. Abundance and density estimate of animals/km² have been calculated from the raw data counts for minke whales. These results are set out below.

11.6.5.5 Review of Abundance and Density Estimates for Minke Whale

155. For the entire SCANS-III survey area, minke whale abundance in the summer of 2016 was estimated to be 13,101 with an overall estimated density of 0.0108/km² (CV = 0.345; 95% CI = 7,050-26,721; **Plate 11-21**, Hammond *et al.* 2021).
156. Within the impact assessments for minke whale, the worst case density estimates from the SCANS-III surveys will be used. The Projects are located in SCANS-III survey block O with a density estimate of (Hammond *et al.* 2021):
- Abundance = 603 minke whale (CV=0.62; 95% CI=109-1,670)
 - Density = 0.0100 minke whale/km² (CV=0.621)




Plate 11-21 Estimated Minke Whale Density in Each SCANS-III Survey Block (Hammond *et al.* 2021)

157. Genetic evidence suggests that the minke whales of the North Atlantic are likely to be a single genetic population (Anderwald *et al.* 2012). Therefore, IAMMWG (2022) considers a single MU is appropriate for minke whales in European waters.
158. The single MU for minke is the CGNS MU (**Plate 11-18**), covering the same geographical area as described for white-beaked dolphin and common dolphin (IAMMWG, 2022). The reference population for minke whales in the CGNS MU is 20,118 animals (CV = 0.18; 95% CI = 14,061 – 28,786; IAMMWG, 2022). This estimate was derived from using the SCANS-III (Hammond *et al.* 2017) and ObSERVE data (Rogan *et al.* 2018). The IAMMWG (2022) note the abundance of minke whales is highly seasonal, with abundance peaking during migration south into waters around the UK for summer.

11.6.5.6 Diet of Minke whale

159. Minke whales feed on a variety of fish species, including herring, cod and haddock. Minke whale feed by engulfing large volumes of prey and water, which they then 'sieve' out of through their baleen plates and swallow their prey whole.
160. A study into the diet of minke whale in the north-eastern Atlantic sampled a total of 210 minke whale for stomach contents from 2000 to 2004, with a total of 37 minke whale samples analysed within the northern North Sea. Within this area, minke whale were found to prey upon a number of different species at the population level, however, 84% of individuals were found to prey upon only one species. Sandeels (56% of total prey by biomass) and mackerel (30% of total prey by biomass) were found to be the most dominant prey species for minke whale in the northern North Sea (Windsland *et al.* 2007).

11.6.6 Grey Seal

11.6.6.1 Desk-Based Review of Grey Seal Presence

161. Grey seals only occur in the North Atlantic, Barents and Baltic Sea with their main concentrations on the east coast of Canada and United States of America and in north-west Europe (SCOS 2020).
162. Approximately 36% of the world's grey seals breed in the UK, and 81% of these breeds at colonies in Scotland with the main concentrations in the Outer Hebrides and in Orkney. There are also breeding colonies in Shetland, on the north and east coasts of mainland Britain and in south-west England and Wales (SCOS 2020).
163. Grey seals are wide ranging and can breed and forage in different areas (Russell *et al.* 2013). Maximum foraging ranges for grey seal has been recorded to be up to 448km (Carter *et al.* 2022). For example, tags deployed on grey seals at Donna Nook and Blakeney Point in May 2015, indicated that they used multiple haul-outs sites; with one hauling out in the Netherlands and one in Northern France (Russell 2016). **Plate 11-22** shows the tagged seal movements along the east coast of England and indicates that grey seal travel between haul-out sites along the east coast of England, as well as to the north of France, Firth of Forth and Dogger Bank, and travel through the DBS project areas (Russell 2016).
164. The north Dutch coastline is also an important foraging zone and migration route for grey seal (Brasseur *et al.* 2010). A study on the grey seal population in the Dutch part of the Wadden Sea shows that the growth of the breeding population is fuelled by the annual immigration of grey seals from the UK, indicating connectivity with the Wadden Sea area (Brasseur *et al.* 2018).

165. This is shown through further telemetry tagging studies of grey seals, undertaken from key haul-out sites along the north coast of France (for tagged individuals from 2012; Vincent *et al.* 2017). The results of this tagging study show connectivity of grey seals from the east coast of England to the north coasts of France, Belgium, and the Netherlands, including the Wadden Sea (**Plate 11-23**).
166. The Wadden Sea covers the coastal areas of the Netherlands, Germany and Denmark (approximately 500km). It is an intertidal zone in the southeastern North Sea with an area of approximately 10,000km².
167. Coordinated aerial, boat and land surveys of the Dutch, German and Danish Wadden Sea grey seal areas including Helgoland (Germany) are aimed at estimating changes in numbers of grey seal in the Wadden Sea area. Annual surveys are conducted in the Wadden Sea, during the moult and breeding season by the Trilateral Seal Expert Group (TSEG). TSEG counts for adult grey seals were conducted by aerial surveys during the moulting period in the spring 2020. Studies show that in moult period, the animals present are not necessarily animals breeding in the Wadden Sea and considerable exchange occurs with the much larger UK population (Brasseur *et al.* 2015; 2017).
168. The most recent counts in the pupping season in November-January of 2021-2022 resulted in a total of 2,214 pups in the whole Wadden Sea and Helgoland, (Schop *et al.* 2022), representing a growth of 15% compared to 2020- 2021 (Brasseur *et al.* 2021). Previous counts have shown the number of grey seal recorded in the Wadden Sea area has been steadily increasing, with a mean annual 10% increase over the past five years, with the most recent count of grey seal in 2021 being of 7,649, (Schop *et al.* 2022).
169. Unger *et al.* (2022) recorded juvenile grey seals travelling from the haul out sites in Helgoland, Germany throughout the North Sea, showing that grey seals travelling in the DBS array areas can be from the Wadden Seas.




Plate 11-22 Tagged Grey Seal Movements Along the East Coast of England (Russell 2016).




Plate 11-23 Grey Seal Telemetry Tags (Shown in Green are the Results from the Tagging of 11 Individuals in 2012, from the Baie de Somme (BDS) Haul-out Site on the North Coast of France. Red Dots Indicate Haul-out Sites, and Red Circles Indicate Breeding Locations.

170. There is a considerable amount of movement of grey seals that occurs (as observed from telemetry data) among the different areas and regional subunits of the North Sea, and no evidence to suggest that grey seals on the North Sea coasts of Denmark, Germany, the Netherlands or France are independent from those in the UK (SCOS 2021).

11.6.6.2 Grey Seal Haul-Out Sites

171. Compared with other times of the year, grey seals in the UK spend longer hauled out during their annual moult (between December and April) and during their breeding season (SCOS 2021).

172. In eastern England, pupping occurs mainly between early November and mid December (SCOS 2020). Pups are typically weaned 17 to 23 days after birth, when they moult their white natal coat, and then remain on the breeding colony for up to two or three weeks before going to sea. Mating occurs at the end of lactation and then adult females depart to sea and provide no further parental care (SCOS 2020).
173. Principal grey seal haul-out sites are included in **Table 11-16** which shows the approximate distance to the closest point of DBS East array area, DBS West array area, and the offshore export cable corridor; in addition with the most recent grey seal count for each haul out location.

Table 11-16 The Most Recent Grey Seal Count at Each of the Nearby Haul-out Sites, and the Distance to DBS

Haul-out site	Distance to DBS East and DBS West	Grey seal count
Holy Island	223km from landfall* 214km from offshore export cable corridor 225km from DBS East array area 254km from DBS West array area	4,251 (2020 mean grey seal count; SCOS 2020)
Farne Islands, (Northumberland)	212km from landfall* 205km from offshore export cable corridor 215km from DBS East array area 246km from DBS West array area	

Haul-out site	Distance to DBS East and DBS West	Grey seal count
Tees	93km from landfall* 120km from offshore export cable corridor 185km from DBS East array area 200km from DBS West array area	30 (2021 mean grey seal count, SCOS,2023)
Ravenscar	52km from landfall* 62km from offshore export cable corridor 140km from DBS East array area 150km from DBS West array area	Both Ravenscar and Filey Brigg are transient sites. 15 grey seal (grey seal count, Yorkshire seal org, 2023)
Filey Brigg	28km from landfall* 35km from offshore export cable corridor 106km from DBS East array area 132km from DBS West array area	
Donna Nook	62km from landfall* 65km from offshore export cable corridor 153km from DBS East array area 151km from DBS West array area	3,897 grey seal (2021 mean grey seal count; SCOS 2023).

Haul-out site	Distance to DBS East and DBS West	Grey seal count
The Wash	108km from landfall* 126km from offshore export cable corridor 197km from DBS East array area 191km from DBS West array area	799 grey seal (2021 mean grey seal count; SCOS 2023).
Blakeney Point National Nature Reserve (NNR)	131km from landfall* 134km from offshore export cable corridor 177km from DBS East array area 166km from DBS West array area	493 grey seal (2021 mean grey seal count; SCOS 2023).
Horsey	185km from landfall* 181km from offshore export cable corridor 198km from DBS East array area 182km from DBS West array area	380 grey seal (2021 mean grey seal count; SCOS 2023)
Scroby Sands	198km from landfall* 198km from offshore export cable corridor 213km from DBS East array area 195km from DBS West array area	1,377 grey seal (2021 mean grey seal count; SCOS 2023).

*Measured in a straight line from landfall site

11.6.6.3 Results from the Site-Specific Surveys for Grey Seal

174. During the first years site specific surveys from March 2021 to February 2022, 11 grey seals were recorded in DBS East array area on seven surveys. At DBS West array area 31 individuals were recorded during eight surveys. In addition to grey seal, there were a number of sightings that was recorded as 'unidentified seal'; in DBS East array area (n=25) and DBS West array area (n=34). Furthermore, marine mammals that were recorded that could not be attributed to a species were recorded as 'marine mammal species', DBS West array area (n=14), some of which could have been grey seals. The number of definite grey seal recordings peaked in the summer months.
175. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), 92 grey seals were recorded, with peak numbers recorded in January. March, June and July.
176. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), a total of 46 grey seals were recorded. In addition, there were 218 unidentified seals recorded which possibly can be attributed to grey seals, along with 210 unidentified small cetacean or seal species. No species abundance or density estimates were undertaken. There were no obvious seasonal peaks.
177. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there were 244 incidental sightings of grey seals, with 244 individuals. There was also a further 19 sightings of unidentified seal species, with 17 individuals plus two dead seals, with which could be possibly attributed to grey seals. No species abundance or density estimates were undertaken. There was no clear seasonal trend of grey seal, however numbers peaked from December 2010 to April 2011 which would be the pupping, breeding and moulting seasons.
178. Density estimates of animals/km² have been calculated from the raw data counts for (i) grey seals and (ii) unidentified seal species and are set out below. These abundance and densities are for the DBS East array area plus 4km buffer (**Table 11-17; Table 11-19**) and DBS West array area plus 4 km buffer (**Table 11-18; Table 11-20**). No correction factor has been used to count for diving seals, therefore the abundance and density estimates are the absolute minimum.

11.6.6.4 Site-Specific Density Estimates for Grey Seal

179. Density estimates of animals/km² have been calculated from the raw data counts for (i) grey seals and (ii) unidentified seal species and are set out below. These abundance and densities are for the DBS East array area plus 4km buffer (**Table 11-17; Table 11-19**) and DBS West array area plus 4 km buffer (**Table 11-18; Table 11-20**). No correction factor has been used to count for diving seals, therefore the abundance and density estimates are the absolute minimum.
180. The abundance estimates of grey seal and unidentified seal species are presented on a graph in **Plate 11-24** for DBS East array area and **Plate 11-25** for DBS West array area. The months with no sightings are not included in the table.

Table 11-17 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km²) of Grey Seal in: a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer

a) DBS East array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Jun-21	2	17	2	43	0.71	0.02
Jul-21	2	17	2	43	0.71	0.02
Aug-21	3	26	3	78	0.58	0.01
Sep-21	1	9	1	26	1.00	0.01
Oct-21	1	8	1	25	1.00	0.01
Jan-22	1	9	1	26	1.00	0.01
Feb-22	1	9	1	26	1.00	0.01
b) DBS East array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Oct-21	1	8	1	25	1.00	0.02
c) DBS East 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Jun-21	2	18	2	44	0.71	0.04
Jul-21	2	17	2	44	0.71	0.04
Aug-21	3	26	3	70	0.58	0.06
Sep-21	1	9	1	26	1.00	0.02
Jan-22	1	9	1	26	1.00	0.02
Feb-22	1	9	1	26	1.00	0.02

Table 11-18 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km²) of Grey Seal in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer

a) DBS West array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	1	9	1	34	1.00	0.01
Jun-21	6	50	17	92	0.41	0.05
Jul-21	8	67	17	126	0.35	0.07
Aug-21	4	33	8	75	0.50	0.04
Oct-21	1	8	1	25	1.00	0.01
Dec-21	3	25	3	59	0.58	0.03
Jan-22	3	25	3	59	0.58	0.03
Feb-22	5	42	8	84	0.45	0.05
b) DBS West array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Jun-21	2	16	2	41	0.71	0.03
Jul-21	7	58	17	117	0.38	0.12
Aug-21	3	25	3	58	0.58	0.05
Oct-21	1	8	1	25	1.00	0.02
Dec-21	1	8	1	25	1.00	0.02
Jan-22	2	17	2	41	0.71	0.03
Feb-22	4	33	8	67	0.50	0.07
c) DBS West 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	1	9	1	26	1.00	0.02
Jun-21	4	34	8	76	0.50	0.08
Jul-21	1	8	1	25	1.00	0.02
Aug-21	1	8	1	25	1.00	0.02
Dec-21	2	17	2	42	0.71	0.04
Jan-22	1	9	1	26	1.00	0.02
Feb-22	1	9	1	26	1.00	0.02

Table 11-19 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km²) of Unidentified Seal Species in a) DBS East Array Area Plus 4km Buffer; b) DBS East Array Area c) 4km Buffer

a) DBS East array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	3	226	3	60	0.58	0.03
May-21	2	17	2	52	0.71	0.02
Jun-21	3	26	3	61	0.58	0.03
Jul-21	4	34	9	69	0.50	0.04
Sep-21	1	9	1	26	1.00	0.01
Oct-21	1	8	1	25	1.00	0.01
Nov-21	1	9	1	36	1.00	0.01
Dec-21	6	51	17	94	0.41	0.06
Jan-22	2	17	2	43	0.71	0.02
Feb-22	2	17	2	43	0.71	0.02
b) DBS East array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	42	0.71	0.03
Jul-21	1	8	1	25	1.00	0.02
Nov-21	1	9	1	27	1.00	0.02
Dec-21	2	17	2	43	0.71	0.03
Jan-22	2	17	2	43	0.71	0.03
Feb-22	1	9	1	25	1.00	0.02
c) DBS East 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Mar-21	2	17	2	26	0.71	0.03
Jul-21	1	8	1	25	1.00	0.02
Nov-21	1	9	1	27	1.00	0.02
Dec-21	2	17	2	43	0.71	0.03
Jan-22	2	17	2	43	0.71	0.03
Feb-22	1	9	1	25	1.00	0.02

Table 11-20 Raw Counts and Abundance and Density Estimates (no. Estimated Individuals per km²) of Unidentified Seal Species in a) DBS West Array Area Plus 4km Buffer; b) DBS West Array Area c) 4km Buffer

a) DBS West array area plus 4km buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	1	9	1	26	1.00	0.01
May-21	9	75	25	133	0.33	0.08
Jun-21	10	83	33	142	0.32	0.09
Jul-21	5	42	8	84	0.45	0.05
Sep-21	1	8	1	25	1.00	0.01
Oct-21	1	8	1	25	1.00	0.01
Dec-21	3	25	3	59	0.58	0.03
Jan-22	3	25	3	67	0.58	0.03
Feb-22	1	8	1	25	1.00	0.01
b) DBS West array area						
Survey date	Raw count	42	Lower CI	Upper CI	Precision	Density
Apr-21						
May-21	6	50	8	108	0.14	0.1
Jun-21	3	25	3	58	0.71	0.05
Jul-21	2	17	2	42	0.38	0.03
Sep-21	1	8	1	24	0.58	0.02
Oct-21	1	8	1	25	1.00	0.02
Dec-21	1	8	1	25	1.00	0.02
Jan-22	2	17	2	50	0.71	0.03
c) DBS West 4km Buffer						
Survey date	Raw count	Abundance	Lower CI	Upper CI	Precision	Density
Apr-21	1	9	1	26	1.00	0.02
May-21	3	25	3	59	0.58	0.06
Jun-21	7	596	17	118	0.38	0.14
Jul-21	3	25	3	59	0.58	0.06
Dec-21	2	17	2	42	0.71	0.04
Jan-22	1	9	1	34	1.00	0.02
Feb-22	1	9	1	26	1.00	0.02

Dogger Bank South Offshore Wind Farms




Plate 11-24 Estimated Abundance of Grey Seal at DBS East Array Area




Plate 11-25 Estimated Abundance of Grey Seal at DBS West Array Area

11.6.6.5 Review of Abundance and Density Estimates for Grey Seal

181. The following sections provide the grey seal at-sea density estimates from grey seal mapping dataset; Carter *et al.* (2022).
182. The relative seals at-sea density maps have also been used to calculate grey seal density estimates for the DBS array areas. The Carter *et al.* (2022) density maps are an update to the Russell *et al.* (2017) mapping and include updated tagging studies. These density maps only include tagging studies from the UK.
183. The resultant density of seals at-sea maps (Carter *et al.* 2022; **Plate 11-26** and **Figure 11-1-1**) differ from the Russell *et al.* (2017) maps, in that they show the relative density of seals in each 5km by 5km grid cell. Each grid cell shows the percentage of the overall seal population within the British Isles, which can then be related to the current best population estimate for each species. This ensures that the relative densities can be updated based on overall population level changes. To calculate a density estimate to be used in assessments from the Carter *et al.* (2020) data, the current at-sea population of each species must be used. A correction factor is also applied to the overall population level to take account of those individuals that are estimated to be on land, and therefore not included in the density mapping.
184. Carter *et al.* (2022) provides habitat-based predictions of at sea distribution for seals around the British Isles. The habitat preference approach predicted distribution maps provide estimates per species, on a 5km by 5km grid, of relative at sea density for seals hauling-out in the British Isles. It is important to note that Carter *et al.* (2022) provides relative density (i.e. percentage of the total at sea population in each grid at any one time), whereas previous usage maps (Russell *et al.* 2017) have presented absolute density (i.e. number of animals within each grid at any one time).
185. The grey seal density estimates for DBS East array area and DBS West array area have been calculated from the seal at sea usage maps (Carter *et al.* 2022) based on the 5km x 5km grids that overlap with the offshore Project areas. The total grey seal population in the British Isles, at sea, is approximately 159,175 individuals, based on the corrected values and most recent haul-out counts for the UK (SCOS 2020). This is the population estimate used with the Carter *et al.* (2022) data to calculate density estimates for DBS East array area and DBS West array areas.
186. Within the impact assessments for grey seal, density estimates for DBS East and DBS West calculated from the seal at sea usage maps (Carter *et al.* 2022) will be used.
187. The mean at sea relative density estimates for these areas have been calculated from Carter *et al.* (2022);

- 0.181 individuals per km² for the DBS East array area;
- 0.260 individuals per km² for the DBS West array area; and
- 0.531 individuals per km² for the Offshore export cable corridor; and
- 0.386 individuals per km² for the total Offshore Development Area.




Plate 11-26 Relative Density of Grey Seals a) Mean With Associated Cell-wise b) Lower and c) Upper 95% Confidence Intervals of the Percentage of the at Sea Population (Excluding Hauled-out Seals) Estimated to be Present in Each 5km x 5km Grid Cell at Any One Time (Carter et al. 2022)

Grey Seal Population Counts

188. Grey seal population trends are assessed from the counts of pups born during the autumn breeding season, when females congregate on land to give birth (SCOS 2020). The pup production estimates are converted to estimates of total population size (1+ aged population) using a mathematical model and projected forward (SCOS 2020).
189. The most recent surveys of the principal grey seal breeding sites in Scotland, Wales, Northern Ireland and south-west England resulted in an estimate of 68,050 pups (95% CI = 60,500-75,100; SCOS 2020). When the pup production estimates are converted to estimates of total population size, there was an estimated 149,700 grey seals in 2019 (approximate 95% CI = 120,000-174,900; SCOS 2020).
190. Based on 2016 and 2018 pup production and projecting the model forward. This is an increase of approximately 1.4% per year between 2012 and 2019 (SCOS 2020).
191. In the southern North Sea, the rates of increase in pup production from 2010 to 2014 by an average 22% per year suggests that there must be some immigration from colonies further north (SCOS, 2019). The colonies in the southern North Sea are still increasing in population size, but the rate has been much lower in the last three years, giving an early indication that they may be reaching carrying capacity (SCOS 2019).
192. The most recent counts of grey seal in the August surveys 2016-2021, estimated that the minimum count of grey seals in the UK was 42,765 (SCOS 2021).
193. In accordance with the agreed approach for other offshore wind farms, and as agreed during the 2nd ETG meeting on the 21st February 2023, the reference population extent for grey seal will incorporate the south-east England and MU, north-east England MU (IAMMWG 2013; SCOS, 2020).
194. The reference population for grey seal is therefore currently based on the following most recent estimates for the:
 - South-east England MU = 8,667 grey seal (SCOS, 2020); and
 - North-east England MU = 6,501 grey seal (SCOS, 2020);
195. Assessments will be done in the context of the nearest MU as well as the wider reference population. As a worst case it is assumed that all seals are from the nearest MU, the south-east England MU, although the more realistic assessment is based on wider reference population which takes into account movement of seals.

196. The reference population for grey seal is therefore currently based on the most recent estimates as shown in **Table 11-21**.

Table 11-21 Grey Seal Count Population Estimates

Population area	Grey seal haul-out count	Source of haul-out count data	Correction factor for seals not available to count	Grey seal total population
SE England MU	8,667	SCOS 2020	0.2515	34,461
NE England MU	6,501	SCOS 2020	0.2515	25,849
Total wider reference population	15,168	-	0.2515	60,310

197. Having an understanding of these movements of seals in the Wadden Sea and their fluctuating dependency on the different areas is key to managing the populations and ensuring their long-term protection (Unger *et al.* 2022).
198. The total reference population for the assessment is 15,168 grey seal. Assessments are in the context of the nearest MU as well as the wider reference population (of 60,310). As a worst case it is assumed that all seals are from the nearest MU, the south-east England MU, although the more realistic assessment is based on wider reference population which takes into account movement of seals.

11.6.6.6 Diet and Foraging of Grey Seal

199. Grey seals will typically forage in the open sea and return regularly to land to haul-out, although they may frequently travel up to 100km between haul-out sites. Foraging trips generally occur within 100km of their haul-out sites, although grey seal can travel up to several hundred kilometres offshore to forage (SCOS, 2020).
200. Individual grey seals based at a specific haul-out site often make repeated trips to the same region offshore, but will occasionally move to a new haul-out site and begin foraging in a new region (SCOS, 2019). Telemetry studies of grey seal in the UK have identified a highly heterogeneous spatial distribution with a small number of offshore 'hot spots' continually utilised (Matthiopoulos *et al.* 2004; Russell *et al.* 2017).
201. Grey seals are generalist feeders, feeding on a wide variety of prey species (SCOS, 2021; Hammond and Grellier 2006). Diet varies seasonally and from region to region (SCOS 2021).

202. In the North Sea, principal prey items are sandeel, whitefish (such as cod, haddock, whiting and ling *Molva molva*) and flatfish (plaice *Pleuronectes platessa*, sole, flounder, and dab *Limanda limanda*) (Hammond and Grellier, 2006). Amongst these, sandeels are typically the predominant prey species.
203. Food requirements depend on the size of the seal and fat content (oiliness) of the prey, but an average consumption estimate of an adult is 4 to 7kg per seal per day depending on the prey species (SCOS 2021).

11.6.7 Harbour seal

11.6.7.1 Desk-Based Review of Harbour Seal Presence

204. Harbour seals have a circumpolar distribution in the Northern Hemisphere and are divided into five sub-species. The population in European waters represents one subspecies *Phoca vitulina vitulina* (SCOS 2021).
205. On the east coast of Britain harbour seal distribution is generally restricted, with concentrations in the major estuaries of the Thames, The Wash and the Moray Firth (SCOS 2021).
206. SMRU, in collaboration with others, has deployed around 344 telemetry tags on harbour seals around the UK between 2001 and 2012. The spatial distributions indicate harbour seals persist in discrete regional populations, display heterogeneous usage, and generally stay within 50km of the coast (Russell and McConnell 2014). Tagged harbour seals were observed to have a more coastal distribution than grey seals and do not travel as far from haul-outs (**Plate 11-27**; Carter *et al.* 2022).




Plate 11-27 Telemetry Tracks by Deployment Region for Harbour Seals Aged One Year or Over (Russell and McConnell 2014).

207. Harbour seals generally make smaller foraging trips than grey seal, typically travelling 40-50km from their haul-out sites to foraging areas (SCOS, 2019). Tracking studies have shown that harbour seals travel 50-100km offshore and can travel 200km between haul-out sites (Lowry *et al.* 2001; Sharples *et al.* 2012). The range of these trips varies depending on the location and surrounding marine habitat. Tagging studies undertaken on harbour seal at The Wash (2003-2005) have shown that this population travels larger distances for their foraging trips than for other harbour seal populations and repeatedly forage between 75km and 120km offshore (average was 80km), with one seal travelling 220km (Sharples *et al.* 2012). The typical and average foraging range for harbour seal is 50-80km (SCOS 2021), with a maximum foraging range of up to 273km (Carter *et al.* 2022).
208. Research has looked at harbour seal movements from the Wadden Sea and the connectivity into the North Sea. Tougaard *et al.* 2003 recorded three harbour seals undertaking long trips, travelling up to three weeks (**Plate 11-28**), therefore harbour seals in the Wadden Sea MU might travel to the DBS array areas.




Plate 11-28 Selected Examples of Foraging Trips. Purple: Elsam02 On an 18-day Trip to the Deeper Parts of the North Sea.. Red: Elsam05 on a 15-day Trip to the German Bight. Blue: Elsam08 on a 10-day Trip to Horns Reef (Tougaard *et al.* 2003)

209. The most recent counts in the Wadden Sea area in the molt season in 2022 resulted in 23,654 harbour seals a 12% decrease since the count in 2021. In 2022, there was a total of 8,514 pups counted which is 22% lower than the count of 10,903 in 2021 which is the lowest count since 2011 (Galatius *et al.* 2022). Since 2012, the harbour seal population has been stable with an annual increase of 5% (Brasseur *et al.* 2018) until recent years. Having an understanding of these movements of seals in the North Sea and the Wadden Sea along with their fluctuating dependency on the different areas is key to managing the populations and ensuring their long-term protection (Unger *et al.* 2022).

210. Harbour seal come ashore in sheltered waters, typically on sandbanks and in estuaries, but also in rocky areas. Harbour seal regularly haul-out on land in a pattern that is often related to the tidal cycle (SCOS 2021). Harbour seal give birth to their pups in June and July and pups can swim almost immediately after birth (SCOS, 2020). Harbour seals moult in August and spend a higher proportion of their time on land during the moult than at other times (SCOS 2021).
211. Principal harbour seal haul-out sites are included in **Table 11-22**, which shows the approximate distance to the closest point of the Offshore development area, and the most recent harbour seal count for each location. These harbour seal haul-out sites are also shown in **Table 11-22**.
212. The Carter *et al.* (2022) density maps include updated tagging studies (**Plate 11-27**). These density maps only include tagging studies from the UK.
213. The harbour seal density estimates for DBS East and DBS West array areas have been calculated from the seal at sea usage maps (Carter *et al.* 2022) based on the 5km x 5km grids that overlap with the Projects. The total harbour seal population in the British Isles, at sea, is approximately 40,600 individuals, based on the corrected values and most recent haul-out counts for the UK (SCOS, 2021). This is the population estimate used with the Carter *et al.* (2022) data to calculate density estimates for DBS East and DBS West array areas.

11.6.7.2 Harbour Seal Haul-Out Sites

Table 11-22 The Most Recent Harbour Seal Count at Each of the Nearby Haul-out Sites, and the Distance to DBS

Haul-out site	Distance to DBS East and DBS West	Harbour seal count
Tees	93km from landfall 120km from offshore export cable corridor 185km from DBS East array area 200km from DBS West array area	86 (2021 harbour seal count; SCOS, 2023)
Ravenscar	52km from landfall 62km from offshore export cable corridor 140km from DBS East array area 150km from DBS West array area	Both Ravenscar and Filey Brigg are transient sites. <10 harbour seal (harbour seal count, Yorkshire seal org, 2023)

Haul-out site	Distance to DBS East and DBS West	Harbour seal count
Filey Brigg	28km from landfall 35km from offshore export cable corridor 106km from DBS East array area 132km from DBS West array area	
Donna Nook	62km from landfall 65km from offshore export cable corridor 153km from DBS East array area 151km from DBS West array area	122 (2021 harbour seal count; SCOS, 2023)
The Wash	108km from landfall 126km from offshore export cable corridor 197km from DBS East array area 191km from DBS West array area	2,667 (2021 harbour seal count; SCOS, 2023)
Blakeney Point NNR	131km from landfall 1134km from export cable corridor 177km from DBS East array area 166km from DBS West array area	181 (mean 2021 harbour seal count; SCOS, 2023).
Horsey	185km from landfall 181km from offshore export cable corridor 198km from DBS East array area 182km from DBS West array area	12 (mean 2021 harbour seal count; SCOS, 2023).
Scroby Sands	198km from landfall 198km from offshore export cable corridor	25 (mean 2021 harbour seal count; SCOS, 2023).

Haul-out site	Distance to DBS East and DBS West	Harbour seal count
	213km from DBS East array area 195km from DBS West array area	

11.6.7.3 Results from the Site-Specific Surveys for Harbour Seals

214. During the site specific digital aerial surveys of both DBS array areas, undertaken from March 2021 to February 2022, no harbour seals were recorded. However, several sightings were recorded as unidentified seals, some of which could be attributed to harbour seals as well as grey seals (**Table 11-19; Table 11-20**).

11.6.7.4 Results from Surveys in the Dogger Bank Area

215. During the Dogger Bank Tranche C site specific video digital aerial surveys (12 surveys undertaken between January 2013 to December 2013), 12 harbour seals were recorded, with peak numbers recorded in January, and Spring.
216. Throughout the Dogger Bank Zone 3 site specific video digital aerial surveys (26 surveys undertaken between April 2010 to May 2012), six harbour seals were recorded. In addition, there were 218 unidentified seals recorded which possibly can be attributed to harbour seals, along with 210 unidentified small cetacean or seal species. No species abundance or density estimates were undertaken.
217. During the ornithological Surveys for the Forewind Round 3 Dogger Bank Offshore Wind Farm (monthly surveys January 2010 to June 2012 (excluding February 2012)), there were 15 incidental sightings of harbour seals, with 15 individuals. There was also a further 18 sightings of unidentified seal species, with 16 individuals plus two dead seals, with which could be possibly attributed to harbour seals. No species abundance or density estimates were undertaken.

11.6.7.5 Site-Specific Density Estimates for Harbour Seal

218. Based on the DBS site specific surveys, no Harbour seal was recorded during the first year of surveys. However density estimates of unidentified seals/km² have been calculated from the raw data counts for unidentified seal species as there is a possibility that these can be harbour seal (**Table 11-19** and **Table 11-20** in **Section 11.6.6.4**). Unidentified seal had a maximum density of 0.06 seals/km² at DBS East in December 2021 and 0.09 seals/km² in June 2021 at DBS West.

11.6.7.6 Review of Abundance and Density Estimates for Harbour Seal

219. The relative seals at-sea density maps (Carter *et al.* 2022; **Plate 11-29** and **Figure 11-1-2**) have been used to calculate harbour seal density estimates for DBS East and DBS West array areas.




Plate 11-29 Relative Density of Harbour Seals a) Mean With Associated Cell-wise b) Lower and c) Upper 95% Confidence Intervals of the Percentage of the at Sea Population (Excluding Hauled-out Seals) Estimated to be Present in Each 5km x 5km Grid Cell at Any One Time.

220. For harbour seal, it has been estimated that at any one time, 83.4% of all harbour seals may be at-sea (Russell *et al.* 2015), and that 72% of all harbour seals would be hauled-out during the population counts (Lonergan *et al.* 2013); therefore, the total harbour seal population in the British Isles, at-sea, is approximately 49,296 individuals (Carter *et al.* 2022).
221. The harbour seal density estimates for the Projects have been calculated from the latest seal at sea maps produced by SMRU (Carter *et al.* 2022), based on the grids that overlap with each area.
222. The following mean at sea density estimates have been used in the assessment:
- 0.0017 individuals per km² for the DBS East array area;
 - 0.0010 individuals per km² for the DBS West array area;
 - 0.0017 individuals per km² for the offshore export cable corridor; and
 - 0.0015 individuals per km² for the total Offshore Development Area.

Harbour Seal Population Counts

223. Harbour seal are counted while they are on land during their August moult, giving a minimum estimate of population size (SCOS 2021). Combining the most recent counts (2016-2019) gives a total of 31,774 counted in the UK. Scaling this by the estimated proportion hauled out (0.72 (95% CI = 0.54-0.88)) produces an estimated total population for the UK in 2019 of 44,100 harbour seal (approximate 95% CI = 36,100-58,800 ; SCOS, 2021).
224. Recent trends in harbour seal populations (over the last two years) indicate that the harbour seal populations in the Wash are in decline. Since 2010, the numbers of harbour seal in the region had been stable. However, the population count in 2019 was 27% lower than the 2012 to 2018 mean count and the initial examination of the 2020 survey produced a similar estimate (SCOS 2021). This represents a fall of approximately 10%-12% per annum over the two years, or around 25% between 2018 and 2019 (SCOS 2021).
225. In accordance with the agreed approach for other offshore wind farms, and as agreed during the 2nd ETG meeting on the 23rd February 2023, the reference population extent for harbour seal will incorporate the south-east England MU (IAMMWG, 2013; SCOS, 2021).
226. These have also been corrected to take account of the number of seals not available to count during the surveys. Approximately 0.72 harbour seals (Lonergan *et al.* 2013) are available to count within the August surveys (i.e. are hauled-out), and therefore this has been used as a correction factor, to derive total harbour seal numbers within the SE England MU.

227. The reference population for harbour seal is therefore currently based on the following most recent estimates as shown in **Table 11-23**.

Table 11-23 Harbour Seal Counts and Population Estimates in the SE England MU

Population area	Harbour seal haul-out count	Harbour seal haul-out count data	Correction factor for seals not available to count	Harbour seal total population
SE England MU	3,494	SCOS, 2021	0.72	4,853

228. The total reference population for the assessment is currently 3,494 for harbour seal. Assessments will be done in the context of the nearest MU which is the SE England MU.

11.6.7.7 Diet and Foraging of Harbour Seal

229. Harbour seal take a wide variety of prey including sandeels, gadoids, herring and sprat, flatfish, and cephalopods. Diet varies seasonally and regionally, prey diversity and diet quality also showed some regional and seasonal variation (SCOS 2021). It is estimated harbour seals eat 3-5 kg per adult seal per day depending on the prey species (SCOS 2021).

References

Anderwald, P., Brandecker, A., Coleman, M., Collins, C., Denniston, H., Haberlin, M. D., Donovan, M. (2012). Displacement responses of a mysticete, an odontocete, and a phocid seal to construction related vessel traffic. *Endangered Species Research*, 21: 231–240.

APEM. (2022). UK Ornithology and Marine Mammal Baseline Aerial Digital Surveys. Annual Report: RWE Site 1 -March 2021 to February 2022.

APEM. (2022). UK Ornithology and Marine Mammal Baseline Aerial Digital Surveys. Annual Report: RWE Site 2 -March 2021 to February 2022.

Arso Civil, M., Quick, N.J., Cheney, B., Pirotta, E., Thompson, P.M. and Hammond, P.S. (2019). Changing distribution of the east coast of Scotland bottlenose dolphin population and the challenges of area-based management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29, pp.178-196.

Aynsley, C.L. (2017). Bottlenose dolphins (*Tursiops truncatus*) in north-east England: A preliminary investigation into a population beyond the southern extreme of its range. MSc Thesis, Newcastle University.

Banguera-Hinestroza, E., Bjørge, A., Reid, R.J., Jepson, P. and Hoelzel, A.R., (2010). The influence of glacial epochs and habitat dependence on the diversity and phylogeography of a coastal dolphin species: *Lagenorhynchus albirostris*. *Conservation Genetics*, 11(5), pp.1823-1836.

BEIS (2022a). UK Offshore Energy Strategic Environmental Assessment 4 (OESEA4) [Online]. Available at: <https://www.gov.uk/government/consultations/uk-offshore-energy-strategic-environmental-assessment-4-oesea4>. [Accessed March 2023]

Berrow, S.D. and Rogan, E. (1995). Stomach contents of harbour porpoises and dolphins in Irish waters. *European Research on Cetaceans*, 9, pp.179-181.

Börjesson, P., Berggren, P. and Ganning, B. (2003). Diet of harbour porpoises in the Kattegat and Skagerrak seas: accounting for individual variation and sample size. *Marine Mammal Science*, 19(1), pp.38-058.

Brasseur S., Carius F., Diederichs B., Galatius A., Jeß A., Körber P., Meise K., Schop J., Siebert U., Teilmann J., Bie Thøstesen C. & Klöpper S. (2021). EG-Marine Mammals grey seal surveys in the Wadden Sea and Helgoland in 2020-2021. Common Wadden Sea Secretariat, Wilhelmshaven, Germany.

Brasseur S., Cremer J., Czeck R., Galatius A., Jeß A., Körber P., Pund R., Siebert U., Teilmann J. & Klöpper S. (2018). TSEG grey seal surveys in the Wadden Sea and Helgoland in 2017-2018. Common Wadden Sea Secretariat, Wilhelmshaven, Germany.

Brasseur, S., van Polanen Petel, T., Aarts, G., Meesters, E., Dijkman, E., and Reijnders, P. (2010). Grey seals (*Halichoerus grypus*) in the Dutch North Sea: population ecology and effects of wind farms. In: we@sea (Ed.), IMARES Report number C137/10. [Online]. Available at: http://www.we-at-sea.org/leden/docs/reports/RL2-2_2005-006_Effect_of_wind_farms_on_grey_seals_in_the_Dutch_North_Sea.pdf [Accessed March 2023].

Brasseur, S.M.J.M., van Polanen Petel, T.D., Gerrodette, T., Meesters, E.H.W.G., Reijnders, P.J.H. and Aarts G. (2015). Rapid recovery of Dutch grey seal colonies fuelled by immigration. *Marine Mammal Science* 31:405-426.

Brasseur, S.M.J.M., Reijnders, P.J.H., Cremer, J., Meesters, E., Kirkwood, R., Jensen, L.F., Jeß, A., Galatius, A., Teilmann, J & Aarts, G. (2017). Echoes from the past: regional variations in recovery within a harbour seal population. *Plos one*

Brophy, J.t., Murphy, S., Rogan, E. (2009). The diet and feeding ecology of the short-beaked common dolphin (*Delphinus delphis*) in the northeast Atlantic. *International Whaling Commission Scientific Committee Paper SC/61/SM*, 14

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. and Thomas, L. (2001). *Introduction to Distance Sampling. Estimating the abundance of biological populations.* University Press, Oxford.

Canning, S.J., Santos, M.B., Reid, R.J., Evans, P.G.H., Sabin, R.C., Bailey, N and Pierce, G.J. (2008). Seasonal distribution of white-beaked dolphin (*Lagenorhynchus albirostris*) in UK water with new information on diet and habitat use. *Journal of the Marine Biological Association of the UK*. 88(6), 1159-1166.

Canty, A., Ripley, B. (2010). boot: bootstrap R (S-Plus) functions. R package version 1.2-42.

Carter, M.I.D., Boehme, L., Cronin, M.A., Duck, C.D., Grecian, W.J., Hastie, G.D., Jessopp, M., Matthiopoulos, J., McConnell, B.J., Miller, D.L., Morris, C.D., Moss, S.E.W., Thompson, D., Thompson, P.M. and Russell, D.J.F. (2022). Sympatric Seals, Satellite Tracking and Protected Areas: Habitat-Based Distribution Estimates for Conservation and Management. *Front. Mar. Sci.* 9:875869. doi: 10.3389/fmars.2022.875869.

Carter, M.I.D., Boehme, L., Duck, C.D., Grecian, W.J., Hastie, G.D., McConnell, B.J., Miller, D.L., Morris, C.D., Moss, S.E.W., Thompson, D., Thompson, P.M. and Russell, D.J.F. (2020). Habitat-based predictions of at-sea distribution for grey and harbour seals in the British

Isles. Sea Mammal Research Unit, University of St Andrews, Report to BEIS, OESEA-16-76/OESEA-17-78

Clarke, M.R., Santos, M.B. and Pierce, G.J. (1998). The importance of cephalopods in the diets of marine mammals and other top predators. ICES CM, 1000, p.8.

Corkeron, P.J., Bryden, M.M. and Hedstrom, K.E. (1990). Feeding by bottlenose dolphins in association with trawling operations in Moreton Bay, Australia. In *The bottlenose dolphin* (pp. 329-336). Academic Press.

Couperus, A.S. (1997). Interactions between Dutch midwater trawl and Atlantic white-sided dolphins (*Lagenorhynchus acutus*) southwest of Ireland. *J. Northw. Atl. Fish. Sci.* 22:209-218.

Cunningham, L., Baxter, J.M., Boyd, I.L., Duck, C.D., Lonergan, M., Moss, S.E. and McConnell, B. (2009). 'Harbour seal movements and haul-out patterns: implications for monitoring and management'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19 398-407.

De Boer, M.N., 2010. Spring distribution and density of minke whale *Balaenoptera acutorostrata* along an offshore bank in the central North Sea. *Marine Ecology Progress Series*, 408, pp.265-274.

DECC (now Department for Business, Energy and Industrial Strategy (BEIS)) (2016), UK Offshore Energy Strategic Environmental Assessment 3 (OESEA3).

Delefosse, M., Rahbek, M.L., Roesen, L. and Clausen, K.T., (2018). Marine mammal sightings around oil and gas installations in the central North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 98(5), pp.993-1001.

Efron, B., Tibshirani, R.J. (1993). *An introduction to the bootstrap*. Chapman & Hall, London.
R Development Core Team (2012). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.Rproject.org>

Evans P.G.H., Anderwald P. and Baines M.E. (1987) UK Cetacean Status Review. Report to English Nature and Countryside Council for Wales pp. 160. Oxford: Sea Watch Foundation

Fontaine, M.C., Tolley, K.A., Siebert, U., Gobert, S., Lepoint, G., Bouqueneau, J.M. and Das, K., (2007). Long-term feeding ecology and habitat use in harbour porpoises *Phocoena* from Scandinavian waters inferred from trace elements and stable isotopes. *BMC Ecology*, 7, p.1

Fontaine, M.L.C., Roland, K., Calves, I., Austerlitz, F., Palstra, F.P., Tolley, K.A., Ryan, S., Ferreira, M., Jauniaux, T., Llavona, A. and Ürk, B.Ö. (2014). Postglacial climate changes and rise of three ecotypes of harbour porpoises, *Phocoena phocoena*, in western Palearctic waters. *Molecular Ecology*, 23, pp.3306-3321.

Fraser F.C. (1946). Report on Cetacea stranded on the British coasts from 1933 to 1937. No. 12. London: British Museum (Natural History).

Galatius A., Brasseur S., Carius F., Jeß A., Meise K., Meyer J., Schop J., Siebert U., Stejskal O., Teilmann J., Thøstesen C. B. (2022). Survey Results of Harbour Seals in the Wadden Sea in 2022. Common Wadden Sea Secretariat, Wilhelmshaven, Germany.

Gilles, A., Peschko, V., Scheidat, M. and Siebert, U. (2012). Survey for small cetaceans over the Dogger Bank and adjacent areas in summer 2011. Document submitted by Germany to 19th ASCOBANS Advisory Committee Meeting in Galway, Ireland, 20-22 March 2012. AC19/Doc.5-08(P). 16pp.

Gilles, A., Viquerat, S., Becker, E. A., Forney, K. A., Geelhoed, S. C. V., Haelters, J., Nabe-Nielsen, J., Scheidat, M., Siebert, U., Sveegaard, S., van Beest, F. M., van Bemmelen, R. and Aarts, G. (2016). Seasonal habitat-based density models for a marine top predator, the harbour porpoise, in a dynamic environment. *Ecosphere* 7(6):e01367. 10.1002/ecs2.1367.

Hammond, P.S. and Grellier, K. (2006). Grey seal diet composition and prey consumption in the North Sea. Final report to Department for Environment Food and Rural Affairs on project MF0319.

Hammond, P.S., Bearzi, G., Bjørge, A., Forney, K., Karczmarski, L., Kasuya, T., Perrin, W.F., Scott, M.D., Wang, J.Y., Wells, R.S. & Wilson, B. (2008). *Delphinus delphis*. The IUCN Red List of Threatened Species 2008: e.T6336A12649851.

Hammond, P.S., Berggren, P., Benke, H., Borchers, D.L., Collet, A., Heide-Jørgensen, M.P., Heimlich, S., Hiby, A.R., Leopold, M.F., Øien, N. (2002). Abundance of harbour porpoises and other cetaceans in the North Sea and adjacent waters. *J. Appl. Ecol.* 39, 361–376.

Hammond, P.S., Lacey, C., Gilles, A., Viquerat, S., Boerjesson, P., Herr, H., Macleod, K., Ridoux, V., Santos, M., Scheidat, M. and Teilmann, J. (2017). Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. Wageningen Marine Research.

Hammond, P.S., Lacey, C., Gilles, A., Viquerat, S., Boerjesson, P., Herr, H., Macleod, K., Ridoux, V., Santos, M.B., Scheidat, M., Teilmann, J., Vingada, J. and Øien, N. (2021). Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. June 2021. [Online]. Available at:

https://synergy.st-andrews.ac.uk/scans3/files/2021/06/SCANS-III_design-based_estimates_final_report_revised_June_2021.pdf. [Accessed March 2023].

Hammond, P.S., Macleod, K., Berggren, P., Borchers, D.L., Burt, M.L., Cañadas, A., Desportes, G., Donovan, G.P., Gilles, A., Gillespie, D., Gordon, J., Hedley, S., Hiby, L., Kuklik, I., Leaper, R., Lehnert, K., Leopold, M., Lovell, P., Øien, N., Paxton, C., Ridoux, V., Rogan, E., Samarra, F., Scheidat, M., Sequeira, M., Siebert, U., Skov, H., Swift, R., Tasker, M.L., Teilmann, J., Van Canneyt, O. and Vázquez, J.A. (2013). Cetacean abundance and distribution in European Atlantic shelf waters to inform conservation and management. *Biological Conservation* 164: 107-122.

Harmer, S.F. (1927). Report of Cetacea stranded on the British Isles from 1913 to 1926. No. 10. London: British Museum (Natural History).

Heinänen, S. and Skov, H. (2015). The identification of discrete and persistent areas of relatively high harbour porpoise density in the wider UK marine area, JNCC Report No. 544 JNCC, Peterborough.

HiDef (2012). High Resolution Video Survey of Dogger Bank Monthly Reports

HM Government (2011). Marine Policy Statement. [Online]. Available at:

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/69322/pb3654-marine-policy-statement-110316.pdf. [Accessed March 2023].

HM Government (2014). The East Inshore and East Offshore Marine Plans

IAMMWG (2013). Management Units for marine mammals in UK waters (June 2013).

IAMMWG 2022. Updated abundance estimates for cetacean Management Units in UK

IJsseldijk, L.L., Brownlow, A., Davison, N.J., Deaville, R., Haelters, J., Keijl, G., Siebert, U. and ten Doeschate, M.T.I. (2018). Spatiotemporal trends in white-beaked dolphin strandings along the North Sea coast from 1991–2017. *Lutra* 61 (1): 153-163: [Online]. Available at:

https://www.uu.nl/sites/default/files/ijsseldijk_et_al._2018._spatiotemporal_analysis_of_white-beaked_dolphin_strandings._lutra_61_002.pdf. [Accessed March 2023]

Ingram, S.N. and Rogan, E. (2002). Identifying critical areas and habitat preferences of bottlenose dolphins *Tursiops truncatus*. *Marine Ecology Progress Series*. 244, pp.247-255.

JNCC (2010a). JNCC guidelines for minimising the risk of injury to marine mammals from using explosives. August 2010.

JNCC (2010b). Statutory nature conservation agency protocol for minimising the risk of injury to marine mammals from piling noise. August 2010.

JNCC (2017). SAC Selection Assessment: Southern North Sea. January, 2017. Joint Nature Conservation Committee, UK. [Online]. Available at: <http://jncc.defra.gov.uk/page-7243>. [Accessed March 2023]

JNCC (2019). Article 17 Habitats Directive Report 2019: Species Conservation Status Assessments 2019. [Online]. Available at: <https://jncc.gov.uk/our-work/article-17-habitats-directive-report-2019-species/#regularly-occurring-species-vertebrate-species-mammals-marine>. [Accessed March 2023].

JNCC, Natural England and CCW (2010). Draft EPS Guidance - The protection of marine European Protected Species from injury and disturbance. Guidance for the marine area in England and Wales and the UK offshore marine area. Joint Nature Conservation Committee, Natural England and Countryside Council for Wales. October 2010.

Johnston, D.W., Westgate, A.J. and Read, A.J. (2005). Effects of fine-scale oceanographic features on the distribution and movements of harbour porpoises *Phocoena* in the Bay of Fundy. *Marine Ecology Progress Series*, 295, pp.279-293.

Kastelein, R. A., Gransier, R., Hoek, L. and Olthuis, J. (2012). Temporary threshold shifts and recovery in a harbor porpoise (*Phocoena phocoena*) after octave-band noise at 4 kHz. *J. Acoust. Soc. Am.* 132, 3525–3537.

Kastelein, R.A., Hardemann, J. and Boer, H. (1997). Food consumption and body weight of harbour porpoises (*Phocoena phocoena*). In *The biology of the harbour porpoise* Read, A.J., Wiepkema, P.R., Nachtigall, P.E (1997). Eds. Woerden, The Netherlands: De Spil Publishers. pp.

Keiper, C.A., Ainley, D.G., Allen, S.G. and Harvey, J.T. (2005). Marine mammal occurrence and ocean climate off central California, 1986 to 1994 and 1997 to 1999. *Marine Ecology Progress Series*, 289, pp.285-306.

Kinze C.C., Addink M., Smeenk C., Hartmann M.G., Richards H.W., Sonntag R.P. and Benke H. (1997). The white-beaked dolphin (*Lagenorhynchus albirostris*) and the white-sided dolphin (*Lagenorhynchus acutus*) in the North and Baltic Seas: review of available information. *Report of the International Whaling Commission* 47, 675–681.

Learmonth, J.A., Macleod, C.D., Santos, M.B., Pierce, G.J., Crick, H.Q.P. and Robinson, R.A. (2006). Potential effects of climate change on marine mammals. *Oceanography and Marine Biology: An Annual Review* 44, 429-462.

Lindeboom, H., Kouwenhoven, H., Bergman, M., Bouma, S., Brasseur, S., Daan, R., Fijn, R., de Haan, D., Dirksen, S., van Hal, R., Hille Ris Lambers, R., ter Hofstede, R., Krijgsveld, K., Leopold, M. and Scheidat, M. (2011). Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. *Environmental Research Letters*, 6(3).

Liret, C. (2001). *Domaine vital, utilisation de l'espace et des ressources : les grands dauphins, Tursiops truncatus, de l'île de Sein*. Thèse de doctorat de l'Université de Bretagne Occidentale, Brest. 155 p.

Liret, C., Creton, P., Evans, P. G. H., Heimlich-Boran, J. R. and Ridoux, V. (1998). English and French coastal Tursiops from Cornwall to the Bay of Biscay, 1996. Photo-identification Catalogue. Project sponsored by Ministère de l'Environnement, France and Sea Watch Foundation, UK.

Loneragan, M., Duck, C., Moss, S., Morris, C. and Thompson, D., (2013). Rescaling of aerial survey data with information from small numbers of telemetry tags to estimate the size of a declining harbour seal population. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23(1), pp.135-144.

Lowry, L.F., Frost, K.J., Hoep, J.M. and DeLong, R.A. (2001). Movements of satellite-tagged subadult and adult harbor seals in Prince William Sound, Alaska. *Marine Mammal Science* 17(4): 835-861.

Matthiopoulos, J., McConnell, B.J., Duck, C. and Fedak, M.A. (2004). Using satellite telemetry and aerial counts to estimate space use by grey seals around the British Isles. *Journal of Applied Ecology*. 41(3):476-491.

McConnell, B., Loneragan, M. and Dietz, R. (2012). *Interactions between seals and offshore wind farms*. The Crown Estate. ISBN: 978-1-906410-34-5.

Meynier, L. (2004). *Food and feeding ecology of the common dolphin, Delphinus delphis, in the Bay of Biscay: Intraspecific dietary variation and food transfer modelling*. Masters thesis, University of Aberdeen, Aberdeen, United Kingdom. 63pp.

Northridge, S.P., Tasker, M.L., Webb, A. and Williams, J.M., (1995). Distribution and relative abundance of harbour porpoises (*Phocoena L.*), white-beaked dolphins (*Lagenorhynchus albirostris* Gray), and minke whales (*Balaenoptera acutorostrata* Lacepède) around the British Isles. *ICES Journal of Marine Science*, 52(1), pp.55-66.

ORCA, Organistaion cetacea (2023). Short beaked common dolphin. [Online]. Available at: ORCA - Short-beaked common dolphin (orcaweb.org.uk). [Accessed March 2023]

Paxton, C.G.M., Scott-Hayward, L., Mackenzie, M., Rexstad, E. and Thomas, L. (2016). Revised Phase III Data Analysis of Joint Cetacean Protocol Data Resources with Advisory Note, JNCC Report 517, ISSN 0963-8091: <http://jncc.defra.gov.uk/page-7201>.

Pusineri, C., Magnin, V., Meynier, L., Spitz, Hassani, S., Ridoux, V. (2007). Food and feeding ecology of the common dolphin (*Delphinus delphis*) in the oceanic northeast Atlantic and comparison with its diet in neritic areas. *Marine mammal science* 23(1):30-47.

Raum-Suryan, K.L. and Harvey, J.T. (1998). Distribution and abundance of and habitat use by harbor porpoise, *Phocoena*, off the northern San Juan Islands, Washington. *Fishery Bulletin*, 96(4), pp.808-822.

Reeves, R., Smeenk, C., Kinze, C.C., Brownell, R.L. Jr and Lien, J. (1999). White-beaked dolphn *Lagenorhynchus albirostris*, Gray 1846. In

Reid, J.B, Evans, P.G.H. and Northridge, S.P. (2003). Atlas of cetacean Distribution in North west European waters. JNCC, Peterborough.

Rogan, E., Breen, P., Mackey, M., Cañadas, A., Scheidat, M., Geelhoed, S. & Jessopp, M. (2018). Aerial surveys of cetaceans and seabirds in Irish waters: Occurrence, distribution and abundance in 2015-2017. Department of Communications, Climate Action & Environment and National Parks and Wildlife Service (NPWS), Department of Culture, Heritage and the Gaeltacht, Dublin, Ireland. 297pp

Russel, N., Hick., A., Warwick-Evans, V., Ratao, S. (2012). Ornithological Survey Monthly Field Report. Foerewind Dogger Bank Wind Farm Project

Russell, D.J., McClintock, B.T., Matthiopoulos, J., Thompson, P.M., Thompson, D., Hammond, P.S., Jones, E.L., MacKenzie, M.L., Moss, S. and McConnell, B.J., (2015). Intrinsic and extrinsic drivers of activity budgets in sympatric grey and harbour seals. *Oikos*, 124(11), pp.1462-1472.

Russell, D.J.F (2016a). Movements of grey seal that haul out on the UK coast of the southern North Sea. Report for the Department of Energy and Climate Change (OESEA-14-47).

Russell, D.J.F, Jones, E.L. and Morris, C.D. (2017). Updated Seal Usage Maps: The Estimated at-sea Distribution of Grey and Harbour Seals. *Scottish Marine and Freshwater Science* Vol 8 No 25, 25pp. DOI: 10.7489/2027-1.

Russell, D.J.F. and McConnell, B.J. (2014). Seal at-sea distribution, movements and behavior. Report to DECC. URN: 14D/085. March 2014 (final revision).

Russell, D.J.F., Brasseur, S.M.J.M., Thompson, D., Hastie, G.D., Janik, V.M., Aarts, G., McClintock, B.T., Matthiopoulos, J., Moss, S.E.W. and McConnell, B. (2014). Marine mammals trace anthropogenic structures at sea. *Current Biology* Vol 24 No 14: R638–R639.

Santos, M.B. and Pierce, G.J. (2003). The diet of harbour porpoise (*Phocoena phocoena*) in the North east Atlantic. *Oceanography and Marine Biology: an Annual Review* 2003, 41, 355–390.

Santos, M.B., Pierce, G.J., Learmonth, J.A., Reid, R.J., Ross, H.M., Patterson, I.A.P., Reid, D.G. and Beare, D. (2004). Variability in the diet of harbor porpoises (*Phocoena phocoena*) in Scottish waters 1992–2003. *Marine Mammal Science*, 20(1), pp.1–27.

Santos, M.B., Pierce, G.J., Reid, R.J., Patterson, I.A.P., Ross, H.M. and Mente, E., (2001). Stomach contents of bottlenose dolphins (*Tursiops truncatus*) in Scottish waters. *Marine Biological Association of the United Kingdom. Journal of the Marine Biological Association of the United Kingdom*, 81(5), p.873.

SCANS-II (2008). Small cetaceans in the European Atlantic and North Sea. Final Report submitted to the European Commission under project LIFE04NAT/GB/000245, SMRU, St Andrews.

Schop J., Abel C., Brasseur S., Galatius A., Jeß A., Meise K., Meyer J., van Neer A., Stejskal O., Siebert U., Teilmann J., Thøstesen C. B. (2022). Grey Seal Numbers in the Wadden Sea and on Helgoland in 2021-2022. Common Wadden Sea Secretariat, Wilhelmshaven, Germany. [Online]. Available at: <https://www.waddensea-worldheritage.org/resources/2021-2022-grey-seal-report>. [Accessed March 2023].

SCOS (2023). August Seal Counts – England | SMRU (st-andrews.ac.uk) [Online]. Available at: August Seal Counts – England | SMRU (st-andrews.ac.uk). [Accessed March 2023].

SCOS (2020). Scientific Advice on Matters Related to the Management of Seal Populations: 2020. [Online]. Available at: <http://www.smrु.st-andrews.ac.uk/research-policy/scos/>. [Accessed March 2023].

SCOS (2021). Scientific Advice on Matters Related to the Management of Seal Populations: 2021. [Online]. Available at: <http://www.smrु.st-andrews.ac.uk/research-policy/scos/>. [Accessed March 2023].

Sea Watch Foundation (2023). Reports of cetacean sightings eastern England: [Online]. Available at: <http://www.seawatchfoundation.org.uk/recent sightings/>. [Accessed March 2023].

Sea Watch Foundation. (2016). Largest sperm whale stranding ever recorded in the North Sea. Available from: Sea Watch Foundation: Largest Sperm Whale Stranding Ever Recorded In The North Sea.

Sharples R.J., Matthiopoulos, J. and Hammond, P.S. (2008). Distribution and movements of harbour seals around the coast of Britain: Outer Hebrides, Shetland, Orkney, the Moray Firth, St Andrews Bay, The Wash and the Thames, Report to DTI July 2008.

Sharples, R.J., Moss, S.E., Patterson, T.A. and Hammond, P.S. (2012). Spatial Variation in Foraging Behaviour of a Marine Top Predator (*Phoca vitulina*) Determined by a Large-Scale Satellite Tagging Program. PLoS ONE 7(5): e37216.

Silva, M. (1999). Diet of common dolphins, *Delphinus delphis*, off the Portuguese continental coast. J. Mar. Biol. Assoc. UK 79:531-540.

Stone, C.J., Webb, a., Barton, C., Ratcliffe, N., Reed, T.C., Tasker M.L., Camphuysen, C.J. and Pienkowski M.W. (1995). An atlas of seabird distribution in north-west European waters. *Joint Nature Conservation Committee Report*, Peterborough

Teilmann, J. and Carstensen, J. (2012). Negative long term effects on harbour porpoises from a large scale offshore wind farm in the Baltic—evidence of slow recovery. *Environmental Research Letters*, 7(4), p.045101.

Teilmann, J., Carstensen, J., Dietz, R., Edrén, S. and Andersen, S. (2006). Final report on aerial monitoring of seals near Nysted Offshore Wind Farm Technical report to Energi E2 A/S. Ministry of the Environment Denmark.

Tees Seals Research Programme. (2019). Monitoring Report No.31 (1989- 2019). Available from: Tees Seals Research Programme (inca.uk.com).

The Independent (2021). Walrus spotted in Northumberland. Available from: Walrus spotted in Northumberland a month after Wally returned to Arctic | The Independent

The Telegraph (2022). Bottlenose dolphins are now permanent residents of Yorkshire's coast. Available from: Bottlenose dolphins are now permanent residents of Yorkshire's coast (telegraph.co.uk)

Thompson, P.M. and Miller, D., (1990). 'Summer foraging activity and movements of radio-tagged common seals (*Phoca vitulina*. L.) in the Moray Firth, Scotland'. *Journal of applied Ecology*, pp.492-501.

Tolley, K.A. and Rosel, P.E., (2006). Population structure and historical demography of eastern North Atlantic harbour porpoises inferred through mtDNA sequences. *Marine Ecology Progress Series*, 327, pp.297-308.

Tollit, D.J., Black, A.D., Thompson, P.M., Mackay, A., Corpe, H.M., Wilson, B., Parijs, S.M., Grellier, K. and Parlane, S. (1998). 'Variations in harbour seal *Phoca vitulina* diet and dive-depths in relation to foraging habitat'. *Journal of Zoology*, 244(2), pp.209-222.

Tougaard, J., Carstensen, J., Wisch, M.S., Teilmann, J., Bech, N., Skov, H. and Henriksen, O.D. (2005). Harbour porpoises on Horns reef—effects of the Horns Reef Wind farm. *Annual Status Report 2004 to Elsam*. NERI, Roskilde (Also available at: www.hornsrev.dk).

Tougaard, J., Ebbesen, I., Tougaard, S., Jensen, T., Teilmann, J. (2003). Satellite tracking of harbour seals on Hors Reed. Use of the Horns Redd wind farm area and the North Sea.

Tynan, C.T., Ainley, D.G., Barth, J.A., Cowles, T.J., Pierce, S.D. and Spear, L.B. (2005). Cetacean distributions relative to ocean processes in the northern California Current System. *Deep Sea Research Part II: Topical studies in Oceanography*, 52(1), pp.145-167.

UK Government (2011). Marine Policy Statement. [Online]. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/69322/pb3654-marine-policy-statement-110316.pdf. [Accessed March 2023].


Unger B., Baltzer J., Brackmann J., Brasseur S., Brüggmann M., Diederichs B., Galatius A., Geelhoed S.C.V., Huus Petersen H., IJsseldijk L.L., Jensen T. K., Jess A., Nachtsheim D., Philipp C., Scheidat M., Schop J., Siebert U., Teilmann J., Thøstesen C.B., van Neer A. (2022). Marine mammals. In: *Wadden Sea Quality Status Report*. Eds.: Kloepper S. et al. Common Wadden Sea Secretariat, Wilhelmshaven, Germany. Last updated: 06.09.2022. [Online]. Available at: qsr.waddensea-worldheritage.org/reports/marine-mammals. [Accessed February 2023].

Vincent, C., Huon, M., Caurant, F., Dabin, W., Deniau, A., Dixneuf, S., Dupuis, L., Elder, J.F., Fremau, M.H., Hassani, S. and Hemon, A. (2017). Grey and harbour seals in France: Distribution at sea, connectivity and trends in abundance at haul out sites. *Deep Sea Research Part II: Topical Studies in Oceanography*, 141, pp.294-305.

Waggitt, J.J., Evans, P.G., Andrade, J., Banks, A.N., Boisseau, O., Bolton, M., Bradbury, G., Brereton, T., Camphuysen, C.J., Durinck, J. and Felce, T. (2019). Distribution maps of

cetacean and seabird populations in the North-East Atlantic. Journal of Applied Ecology, 57(2), pp.253-269.

WWT (2009). Distributions of Cetaceans, Seals, Turtles, Sharks and Ocean Sunfish recorded from Aerial Surveys 2001-2008. The Wildfowl and Wetlands Trust



**RWE Renewables UK Dogger
Bank South (West) Limited**

**RWE Renewables UK Dogger
Bank South (East) Limited**

**Windmill Hill Business Park
Whitehill Way
Swindon
Wiltshire, SN5 6PB**

